



Tansley review

Functional responses of plants to elevated atmospheric CO₂ – do photosynthetic and productivity data from FACE experiments support early predictions?

Author for correspondence:

Robert S. Nowak

Tel: +1 775 7841656

Fax: +1 775 7844789

Email: nowak@cabnr.unr.edu

Received: 18 July 2003

Accepted: 19 December 2003

doi: 10.1111/j.1469-8137.2004.01033.x

Robert S. Nowak¹, David S. Ellsworth² and Stanley D. Smith³

¹Department of Natural Resources & Environmental Science, University of Nevada–Reno, Reno, NV 89557 USA; ²School of Natural Resources and Environment, University of Michigan, Ann Arbor, MI 48109 USA; ³Department of Biological Sciences, University of Nevada–Las Vegas, Las Vegas, NV 89154 USA

Contents

Summary	253	V. Primary productivity	265
I. Introduction	254	VI. Response of plant functional types	272
II. Early assessments of [CO ₂] responses in natural ecosystems	254	VII. Conclusions	275
III. Global network of FACE sites	256	Acknowledgements	276
IV. Assimilation and leaf N-content	257	References	276

Summary

Key words: net assimilation, photosynthetic downregulation, primary production, bog, forest, grassland, desert, plant functional type.

Results from 16 free-air CO₂ enrichment (FACE) sites representing four different global vegetation types indicate that only some early predictions of the effects of increasing CO₂ concentration (elevated [CO₂]) on plant and ecosystem processes are well supported. Predictions for leaf CO₂ assimilation (A_{net}) generally fit our understanding of limitations to photosynthesis, and the FACE experiments indicate concurrent enhancement of photosynthesis and of partial downregulation. In addition, most herbaceous species had reduced leaf nitrogen (N)-content under elevated [CO₂] and thus only a modest enhancement of A_{net} , whereas most woody species had little change in leaf N with elevated [CO₂] but a larger enhancement of A_{net} . Early predictions for primary production are more mixed. Predictions that enhancement of productivity would be greater in drier ecosystems or in drier years has only limited support. Furthermore, differences in productivity enhancements among six plant functional types were not significant. By contrast, increases in productivity enhancements with increased N availability are well supported by the FACE results. Thus, neither a resource-based conceptual model nor a plant functional type conceptual model is exclusively supported by FACE results, but rather both species identity and resource

availability are important factors influencing the response of ecosystems to elevated $[\text{CO}_2]$.

© *New Phytologist* (2004) **162**: 253–280

I. Introduction

The increase in atmospheric CO_2 concentration ($[\text{CO}_2]$) during the last 250 yr is unequivocal, and atmospheric $[\text{CO}_2]$ will continue to increase at least for the next several decades. The direct measurements of atmospheric $[\text{CO}_2]$ at Mauna Loa, Hawaii by Keeling and colleagues clearly show a $55 \mu\text{mol mol}^{-1}$ increase since 1959 (Keeling & Whorf, 2002). Longer-term data indicate an atmospheric $[\text{CO}_2]$ near $280 \mu\text{mol mol}^{-1}$ before 1750 (Houghton *et al.*, 2001). Thus, the atmospheric $[\text{CO}_2]$ of $370 \mu\text{mol mol}^{-1}$ in 2001 represents a 32% increase in the last 250 yr, with more than half of that increase occurring in the last 42 yr. Although predictions of future atmospheric $[\text{CO}_2]$ require a number of assumptions about economic growth, technological advances, and carbon sequestration by biological and geological processes, estimates of atmospheric $[\text{CO}_2]$ in the year 2100 range between $540 \mu\text{mol mol}^{-1}$ and $970 \mu\text{mol mol}^{-1}$ (Houghton *et al.*, 2001). Note that for convenience and familiarity, we use concentration units (i.e. $\mu\text{mol mol}^{-1}$) rather than partial pressures (i.e. Pa) to express the content of CO_2 in air, recognizing that changes in elevation among sites, and hence differences in total atmospheric pressure, require appropriate adjustments to the concentration units for physiological processes such as photosynthesis.

Elevated atmospheric $[\text{CO}_2]$ has interested biological scientists, especially ecologists and plant physiologists, because of the potential biological impacts from CO_2 -induced global warming and from direct effects of elevated $[\text{CO}_2]$ on vegetation that are independent of global warming. This latter interest is the focus of this review, and readers interested in the effects of global warming on biological systems are referred to Watson *et al.* (1998) and Melillo *et al.* (2001). The current review augments previous reviews (Ceulemans & Mousseau, 1994; Medlyn *et al.*, 2001) and special issues of *New Phytologist* (Volume 147, Issue 1, 2000; Volume 150, Issue 2, 2001) on the effects of elevated $[\text{CO}_2]$ on specific plants or ecosystems. Our review differs from these earlier papers in that it focuses on whole-ecosystem free-air CO_2 enrichment (FACE) experiments, which currently represent the best technology to manipulate $[\text{CO}_2]$ for ecosystems under natural conditions, and explicitly contrasts the results from these FACE experiments in the context of early predictions made *c.* 20 yr ago on how elevated $[\text{CO}_2]$ may affect different plants and ecosystems.

II. Early assessments of $[\text{CO}_2]$ responses in natural ecosystems

Although elevated $[\text{CO}_2]$ has long been known to stimulate plant growth, particularly within the field of horticulture, the

realization that increasing $[\text{CO}_2]$ would have potentially important effects on crops, forests, and natural ecosystems did not generate interest until the later part of the twentieth century. An early set of reviews on the subject (Kramer, 1981; Kimball, 1983; Percy & Björkman, 1983; Strain & Bazzaz, 1983; Cure & Acock, 1986; Strain, 1987) set the stage for a strong emphasis on elevated $[\text{CO}_2]$ research in the ecological sciences community. The consensus of these early reviews was that increases in $[\text{CO}_2]$ will have direct and relatively immediate effects on two physiological processes in plants: (1) it increases the rate of photosynthetic CO_2 assimilation; and (2) it decreases stomatal conductance. In combination, these effects were thought to substantially increase intrinsic water-use efficiency (WUE; carbon gain per unit water loss) in plants (Morison, 1985). Studies also indicated that plants may produce tissues with lower nitrogen (N) concentration, although it was not known if that was due to a reallocation of N out of leaves or to a dilution effect associated with increased carbohydrate production. These early reviews were also unanimous in articulating great uncertainty about the long-term and whole-ecosystem responses to elevated atmospheric $[\text{CO}_2]$ due to a lack of appropriate data and experiments.

Based on these fundamental observations, a conceptual model proposed by Strain & Bazzaz (1983) predicted increased plant growth and ecosystem productivity in response to elevated $[\text{CO}_2]$ as water availability decreases or nutrient availability increases. Therefore, ecosystems with ample water and low nutrient availability (e.g. bogs and tropical forests) should be relatively unresponsive to elevated $[\text{CO}_2]$; an intermediate response would be observed in systems that are either both water- and nutrient-limited (e.g. upland desert soils and serpentine or calcareous grasslands) or have ample amounts of both water and nutrients (some forests and mesic grasslands); and ecosystems that should respond most strongly would be water-limited yet have adequate nutrients (e.g. alluvial desert soils, fertile dry grasslands). Strain & Bazzaz (1983) thus proposed a testable set of hypotheses that would help guide research on the effects of elevated $[\text{CO}_2]$ on plant productivity. These early reviews, citing our uncertainty of plant responses to elevated $[\text{CO}_2]$ under a variety of ecological conditions, stimulated a huge amount of research on the physiological responses of plants to elevated $[\text{CO}_2]$. For example, the WUE hypothesis, in particular, was the basis for predictive models that simulated changes in global primary production (Melillo *et al.*, 1993) and potential shifts in species composition (Neilson, 1995) as a function of increasing $[\text{CO}_2]$.

An area of particular interest to global change scientists has been to derive an estimate of net primary production (NPP) that would (1) widely accommodate differences in vegetation

type, climate and soil nutrient availability, and (2) be useful in regional and global carbon cycle models. One important approach has been the β factor, or 'biotic growth factor' (Bacastow & Keeling, 1973). The β factor has been variously defined as an empirical coefficient that relates a logarithmic response of NPP to increasing $[\text{CO}_2]$ (Amthor & Koch, 1996) or as a fractional change in NPP with a fractional change in $[\text{CO}_2]$ (Luo & Mooney, 1996). Bacastow & Keeling (1973) gave an initial estimate of β as a likely range of 0.2–0.6 for a recent 10-year increase in $[\text{CO}_2]$. A variety of estimates of the β factor have been put forward, but perhaps the most extensive estimate to date – based on 398 observations from 73 forest tree species – yielded a β factor of 0.43, which translates into an increase in NPP of 32% in response to a doubling (to $700 \mu\text{mol mol}^{-1}$) of $[\text{CO}_2]$ (Wullschleger *et al.*, 1997). Based on calculations in Amthor & Koch (1996), a β of 0.43 would yield an approximate 20% increase in NPP in response to an increase in $[\text{CO}_2]$ to $550 \mu\text{mol mol}^{-1}$, the operational set-point of the global FACE network. However, Luo & Mooney (1996) cautioned that there are 'a very wide range of β -values between, and even within, ecosystems'.

Given the potential for strong variation in plant growth to elevated $[\text{CO}_2]$ across species and ecosystems, it is not surprising that our predictive capability for ecosystem-level responses such as NPP is much less than for individual physiological responses. Early reviews of ecosystem responses to elevated $[\text{CO}_2]$ (Strain, 1987; Bazzaz, 1990; Mooney & Koch, 1994) emphasized this difficulty, and called for larger-scale experiments that specifically examined ecosystem responses to changes in $[\text{CO}_2]$ and other global change phenomena. The need for these experiments was further emphasized by evidence of important feedback mechanisms between below-ground carbon inputs and soil microbial processes (Díaz *et al.*, 1993; Zak *et al.*, 1993) and evidence of novel responses in model terrestrial ecosystems (Jones & Thompson, 1998). These considerations, among others, led Körner (2000) to conclude that an ecosystem approach, which uses fully coupled plant–soil systems and considers potential nonlinear responses, was essential if we are to make meaningful predictions about how an integrated biosphere will respond to global change.

A big question has been: How does the ecological research community reach that goal? Although a modeling approach is one way of trying to elucidate the response of complex, interrelated systems to global change, the information base to parameterize models is much more extensive when parameterizing gas exchange responses to elevated $[\text{CO}_2]$ than it is for other processes that have important ecosystem-level ramifications, such as carbon partitioning and nutrient uptake (Luo *et al.*, 1999). This disparity in available information is largely a consequence of the methodological history of elevated $[\text{CO}_2]$ research. Most early investigations were conducted in controlled environments or glasshouses with herbaceous plants and tree seedlings in pots. These studies had two important limitations (among others): (1) limited rooting volume, and

therefore artificial growing conditions, that could influence the $[\text{CO}_2]$ response of plants (McConnaughay *et al.*, 1993); and (2) the inability to scale seedling responses to whole trees and forest stands (Strain, 1987; Körner, 1995). This led to an extensive network of open-top chamber (OTC) studies that could be conducted with plants in their natural environment. Although these experiments were constrained by (1) the well known 'chamber effect', in which the chamber itself significantly altered microclimate around the plots being investigated (Leadley & Drake, 1993) and (2) the inability to enclose large plots, they provided valuable information, particularly in short-stature herbaceous communities such as a salt marsh (Drake, 1992) and grassland (Owensby *et al.*, 1999). A number of OTC studies were also conducted with forest seedlings and saplings, and although these studies provided important data on tree responses to elevated $[\text{CO}_2]$, they were unable to scale results to closed-canopy forests because they were measuring small trees in the exponential phase of growth (Norby *et al.*, 1999). These limitations with controlled-environment and OTC studies led to the establishment of a new experimental program that utilized free-air CO_2 enrichment (FACE) technology. FACE technology delivers consistent and uniform elevated $[\text{CO}_2]$ treatments to large plots of an intact ecosystem without walls and has two key advantages over other technologies. First, FACE experiments often have plot diameters of 25–30 m, a size large enough to encompass the small-scale spatial structure of the ecosystem. Second, alterations of microclimate are minimal when the FACE system is operated properly to avoid night-time temperature effects (Pinter *et al.*, 2000), and disturbance of natural soil and root processes is avoided (Allen, 1992; Hendrey & Kimball, 1994; Hendrey *et al.*, 1999). Thus, FACE systems allow us to examine complex natural ecosystem responses to elevated $[\text{CO}_2]$ in the field (McLeod & Long, 1999), particularly with regard to temporal and spatial variations in the natural environment (Saxe *et al.*, 1998; Norby *et al.*, 2001).

Because results from FACE systems are likely to capture representative responses of ecosystems to elevated $[\text{CO}_2]$ and given that results from chamber studies have been subject to extensive reviews previously (e.g. Curtis, 1996; Saxe *et al.*, 1998; Medlyn *et al.*, 1999), results from the global network of FACE sites form the basis for this review. We also focus primarily on noncrop systems; Kimball *et al.* (2002) recently summarized results for agricultural crop systems. Our objectives were twofold: (1) to examine whether FACE data support early predictions of plant and ecosystem responses to increasing atmospheric $[\text{CO}_2]$ (specifically, responses of assimilation, leaf N content, primary productivity, and functional groups); and (2) to examine if a rational ecological framework accounts for the observed responses. Specific hypotheses, based on early predictions in the literature, were that: (1) acclimatization (i.e. downregulation) of photosynthesis would occur most prevalently in ecological situations where N is limiting; (2) the response of productivity to elevated $[\text{CO}_2]$ would be greater in drier ecosystems as well as in dry years within more humid environments; (3) based on




Fig. 1 Geographic distribution of noncrop free-air CO₂ enrichment (FACE) sites with labels indicating the type of global ecosystem represented. Grassland sites include native grasslands and seminative meadows and pastures. Sites whose results are used in this review are shaded grey.




Fig. 2 Mean annual temperature and precipitation of noncrop free-air CO₂ enrichment (FACE) sites, except for the bog (BERI) sites. Closed symbols are sites whose results are used in this review. Triangles, forest FACE sites; circles, grassland sites; inverted triangle, savannah; diamond, desert. Global ecosystem categories and their climatic ranges are after Whittaker (1970).

previous estimates of the β factor, NPP at FACE sites should vary around a mean increase of 20% at 550 $\mu\text{mol mol}^{-1}$ [CO₂]; and (4) functional groups such as legumes, C₃ plants, and herbaceous perennials should be more responsive to elevated [CO₂] than woody perennials or C₄ plants.

III. Global network of FACE sites

A total of 24 noncrop FACE sites are situated around the world (Fig. 1). The majority of sites are in Europe, and the

majority of the European sites are organized into two multisite networks: four bog sites in the Bog Ecosystem Research Initiative (BERI) and five grassland sites in a project entitled 'Managing European Grasslands as a Sustainable Resource in a Changing Climate' (MEGARICH). Although five different types of global ecosystems are represented in this network (bogs plus the four shown in Fig. 2), 75% of the current sites are either temperate forest or grassland vegetation. Five global ecosystems do not have any FACE sites, although some (e.g. tundra and tropical forests) have been studied in controlled-environment

facilities or *in situ* using OTCs. Only one site is in a tropical climate (mean annual temperature > 18°C), and no sites receive, on average, > 1500 mm of precipitation each year.

For most sites, CO₂ enrichment occurs using either: (1) the Brookhaven National Laboratory design (Lewin *et al.*, 1994; Hendrey *et al.*, 1999), where blowers mix ambient air with pure CO₂ and then distributes the enriched air across the plot through a series of stand pipes (eight sites); or (2) a pure-CO₂ injection technique (Miglietta *et al.*, 2001a, 2001b; Pepin & Körner, 2002), where pure CO₂ is injected through holes under pressure and turbulent mixing distributes the enriched air across the plot (10 sites). Twenty-one sites have an effective [CO₂] within 10% of 550 µmol mol⁻¹ during daylight hours; 19 of these control for a set [CO₂], whereas the other two maintain an 'ambient plus 200 µmol mol⁻¹' [CO₂]. Of the remaining sites, two sites control for [CO₂] < 495 µmol mol⁻¹ and one site controls for *c.* 700 µmol mol⁻¹. It is important to note that the long-term, integrated [CO₂] achieved by these experiments always is less than their target [CO₂] because of programmed and unprogrammed periods of time when the [CO₂] control is off. Furthermore, ambient [CO₂] also varies among sites, especially at night and in the early morning when plant respiration increases local [CO₂]. The Carbon Dioxide Information Analysis Center (<http://cdiac.esd.ornl.gov>) provides links to individual FACE web sites as well as other useful web sites.

For this review, we primarily extracted results from journal articles that pertain to an individual FACE site. Unfortunately, published data were not available from all sites – the sites used in this review are indicated in Fig. 1 (shaded names). Table 1 lists general information and literature citations on each site whose results we analysed in this review. Although most sites focus solely on [CO₂] treatments, some sites have multifactor experimental designs; for example, complete factorial designs with [CO₂] and one or more other global change factors such as N, precipitation, ozone, or biodiversity. While such interactions are of great interest and offer powerful insights into how other global change factors may modify [CO₂] responses, we did not want to confound the [CO₂] effects with these other factors. Thus, we have generally limited our use of data to only those experimental factors that had all ambient conditions with and without elevated [CO₂]; exceptions are noted in Table 1 and the text. In addition, three sites had two different experiments: ETH-Z and SCG had a managed grassland experiment and a second experiment that examined responses of different plant functional types, and FACTS I has an unreplicated prototype experiment that began 3 yr before the replicated main experiment. Results from both experiments are used as appropriate and are treated as independent studies.

Some precautionary thoughts on our data sets follow. First, data are limited in a number of ways, such as coverage of global ecosystems and climates, number of years of operation, and types of data available. These limitations reflect the nature of

the global FACE network: the great expense to build and operate a FACE site generally limits them to developed countries, which are primarily in temperate ecosystems, and even these developed countries do not always adequately fund the sites. These limitations also highlight the importance of maintaining the current infrastructure and allocating adequate resources to FACE experiments, especially multifactorial experiments and experiments in nontemperate or extreme climates. Second, individual data points within any of our data sets are not true experimental replicates because all other factors that may affect the vegetation responses (e.g. weather, soil characteristics, vegetation management such as grazing or cutting, etc.) are not the same among all FACE sites. We have partly compensated for this variation among sites by expressing most data as the ratio of the response under elevated [CO₂] to that under ambient [CO₂] (i.e. elevated/ambient or E/A), although this approach is not without limitations (Jasienski & Bazzaz, 1999). Furthermore, given that we expect a greater random error for statistical analyses of the data (Filion *et al.*, 2000), we have also carefully examined statistical results when the *P*-value was between 0.05 and 0.10. Although we typically use '*P* ≤ 0.05' as the standard for statistical significance, we also emphasize meaningful results (with the associated *P*-value) when 0.05 < *P* ≤ 0.10. Third, measurement protocols differed among sites, and in some cases, even among years for a particular site. Again, these differences among protocols potentially confound statistical comparisons of data. To the extent possible, we used results that yielded the most comparable measures of vegetation responses. For example, measurement protocols to determine above-ground production in forests (i.e. allometric measurements of tree size) differ from those in grasslands (i.e. destructive harvests at set time intervals), but both protocols result in comparable measures of above-ground production in units of g m⁻² yr⁻¹. Fourth is that the actual [CO₂] treatment varies among sites, as discussed above. Finally, our goal in this review was to examine broad patterns of plant and ecosystem [CO₂] responses to elevated [CO₂] across different resource levels characterized by different sites and not necessarily to characterize the response of a specific flora. Thus, many interesting and important results that pertain to an individual site are not discussed here – the literature cited throughout the text as well as listed in Table 1 provide detailed information on particular sites.

IV. Assimilation and leaf N content

The biochemical basis of leaf CO₂ assimilation responses to short-term increases in [CO₂] is well-established (Farquhar *et al.*, 1980). [CO₂] below 600 µmol mol⁻¹ is typically insufficient to saturate carboxylation in photosynthesis (Fig. 3). Thus, short-term increases in CO₂ supply at [CO₂] below 600 µmol mol⁻¹ tend to increase leaf CO₂ assimilation (*A*_{net}) because carboxylation by Rubisco increases due to increased substrate availability and the suppression of competitive

Table 1 List of free-air CO₂ enrichment (FACE) sites whose data we analysed in this review, along with some site and operating characteristics, sources for data, and explanatory notes for the sites or data sets

Site name and abbreviation used in text	Vegetation type	Country: latitude, longitude	FACE design ¹	Target [CO ₂] ²	Source for data ³	Notes
Bog Ecosystem Research Initiative (BERI)	Bog	Finland ⁴ , Sweden, Netherlands, Switzerland	Pure CO ₂	560 (24 h; season)	<i>Production</i> : for specific species – Heijmans <i>et al.</i> (2001) Table 2; for ecosystem – Hoosbeek <i>et al.</i> (2001) Tables 3 and 4	<i>Production</i> : Results are after 3 yr of elevated [CO ₂]
Oak Ridge National Laboratory: Sweetgum plantation (ORNL)	Forest (deciduous)	USA: 35°54' N, 84°20' W	BNL	565 (24 h, season)	<i>Assimilation</i> : Gunderson <i>et al.</i> (2002) Table 1, Figs 2 and 3 <i>Precipitation</i> : NCDC Oak Ridge ATDD (406750) <i>Production</i> : Belote <i>et al.</i> (2003) Table 1; Norby <i>et al.</i> (2002) Table 1; R. J. Norby (unpublished).	<i>Production</i> : Results are for sweetgum (<i>Liquidambar styraciflua</i>) only; understory < 6% NPP
Forest atmosphere carbon transfer and storage I: Loblolly pine (FACTS I)	Forest (conifer)	USA: 35°59' N, 79°6' W	BNL	Ambient + 200 (24 h, 365 d)	<i>Assimilation</i> : Herrick & Thomas (2001) Fig. 2; Rogers & Ellsworth (2002) Fig. 2 <i>Precipitation</i> : Schäfer <i>et al.</i> (2002) Table 5; NCDC Chapel Hill 2 W (311677) <i>Production</i> : Full experiment – DeLucia <i>et al.</i> (1999) Table 2; Finzi <i>et al.</i> (2002) Table 1; D. Moore & E. H. DeLucia (unpublished); Matamala & Schlesinger (2001) Table 1; Pritchard <i>et al.</i> (2001) Table 3. Prototype – Oren <i>et al.</i> (2001) Fig. 1	Results from both the unreplicated prototype ring and full experiment are used. Prototype APP is woody biomass only. Prototype Target [CO ₂] = 550 (d, season)
Poplar plantation (PopFACE)	Forest (deciduous)	Italy: 42°22' N, 11°48' E	Pure CO ₂	550 (d, season)	<i>Assimilation</i> : Bernacchi <i>et al.</i> (2003) <i>Production</i> : Gielen <i>et al.</i> (2001) Fig. 7	Plots were irrigated; thus precipitation data were not relevant <i>Production</i> : Used LAI as index for APP, averaged over three poplar varieties
Forest atmosphere carbon transfer and storage II: Aspen plantation (FACTS II)	Forest (deciduous)	USA: 45°36' N, 89°42' W	BNL	Ambient + 200 (d, season)	<i>Assimilation</i> : Takeuchi <i>et al.</i> (2001) Table 3 <i>Precipitation</i> : NCDC Rhinelander (477113) <i>Production</i> : Isebrands <i>et al.</i> (2001) Table 4; King <i>et al.</i> (2001) Fig. 1; Percy <i>et al.</i> (2002) Fig. 1; D. F. Karnosky & M. E. Kubiske (unpublished).	<i>Production</i> : Used yearly increment of wood volume as index of APP. Used live + dead root standing crop after 2 yr elevated [CO ₂] as index of BPP

Swiss alpine treeline (SAT)	Forest (conifer)	Switzerland: 46°45' N, 9°45' E	Pure CO ₂	550 (d, season)	<i>Precipitation:</i> Hättenschwiler <i>et al.</i> (2002) Table 1 <i>Production:</i> Hättenschwiler <i>et al.</i> (2002) Fig. 6	<i>Production:</i> Used shoot length, averaged over lead and lateral shoots and over two species, as index of APP
Swiss Eschikon grassland (ETH-Z)	Grassland (perennial)	Switzerland: 47°27' N, 8°41' E	BNL	600 (d, season)	<i>Assimilation:</i> Ainsworth <i>et al.</i> (2003) Fig. 5 <i>Precipitation:</i> Hebeisen <i>et al.</i> (1997) Table 1; Daepp <i>et al.</i> (2000) Table 1; WMO Zurich (646066600) <i>Production:</i> Managed grassland experiment – Hebeisen <i>et al.</i> (1997) Figs 1, 3 and 4; Daepp <i>et al.</i> (2000) Fig. 1. Diversity experiment – Lüscher <i>et al.</i> (1998) Table 3	Results from both experiments (managed grassland and functional type) are used. <i>Production:</i> Excluded partial fumigation year (1993). Managed grassland experiment – averaged over different cutting treatments; averaged monoculture and mixture results if both available; only the low N treatment was used unless otherwise noted
Swiss calcareous grassland (SCG)	Grassland (perennial)	Switzerland: 47°33' N, 7°34' E	Screen Aided CO ₂ Control	600 (24 h, season)	<i>Precipitation:</i> Niklaus <i>et al.</i> (2001) Table 1 <i>Production:</i> Managed grassland experiment – Leadley <i>et al.</i> (1999) Figs 2 and 3. Diversity experiment – Niklaus <i>et al.</i> (2001) Figs 1 and 3	Results from both experiments (managed grassland and biodiversity) are used
Irish seminatural grassland (MR-Irish)	Grassland (perennial)	Ireland: 52°48' N, 6°54' W	Pure CO ₂	600 (d, season)	<i>Production:</i> Bryne & Jones (2002) Table 5	
Grazed pasture grassland (NZGraze)	Grassland (perennial)	New Zealand: 40°14' S, 175°16' E	BNL	475 (d, 365 d)	<i>Assimilation:</i> von Caemmerer <i>et al.</i> (2001) Fig. 5 <i>Precipitation:</i> P. C. D. Newton (unpublished) <i>Production:</i> Morgan <i>et al.</i> (2004b) Fig. 2; P. C. D. Newton (unpublished)	
Biodiversity, CO ₂ , and Nitrogen experiment (BioCON)	Grassland (perennial)	USA: 45°24' N, 93°12' W	BNL	550 (d, season)	<i>Assimilation:</i> Lee <i>et al.</i> (2001) Table 1; Lee <i>et al.</i> (2003) Fig. 2 <i>Precipitation:</i> NCDL Cedar (211390) <i>Production:</i> Reich <i>et al.</i> (2001a) Fig. 2; Reich <i>et al.</i> (2001b) Table 1; D. S. Ellsworth (unpublished)	Multi-factor experiment <i>Production:</i> Results averaged over 2 yr and diversity levels except for functional type results, which were for the low N treatment only

Table 1 continued

Site name and abbreviation used in text	Vegetation type	Country: latitude, longitude	FACE design ¹	Target [CO ₂] ²	Source for data ³	Notes
Jasper Ridge Global Change Experiment (JRGCE)	Grassland (annual)	USA: 37°24' N, 122°14' W	Pure CO ₂	700 (24 h, season)	<i>Precipitation</i> : J. S. Dukes (unpublished) <i>Production</i> : J. S. Dukes (unpublished); Shaw <i>et al.</i> (2002) Table 1; Zavaleta (2001) Chapter 5, Fig. 13	Multi-factor experiment <i>Production</i> : Results from only all-ambient and all-ambient except CO ₂ treatments are used except for comparisons of functional types, which averaged over both water and N treatments
Nevada Desert FACE Facility (NDFE)	Desert	USA: 36°39' N, 115°55' W	BNL	550 (24 h, 365 d)	<i>Assimilation</i> : Hamerlynck <i>et al.</i> (2000) Table 1, Fig. 1; Housman (2002); Naumburg <i>et al.</i> (2003) Fig. 1 <i>Precipitation</i> : http://www.unlv.edu/Climate_Change_Research/Data_Bases/data_index.htm <i>Production</i> : Housman (2002); DeFalco (2003); D. L. Phillips (unpublished); Smith <i>et al.</i> (2000) Figs 1 and 2; S. F. Zitzer (unpublished)	<i>Production</i> : As an index of APP in a year, shoot production for three shrub and two perennial grass species was first weighted by plant cover. Then perennial shoot production was averaged with total production of the four dominant annual species using a 2 : 1 weighting to estimate total above-ground production. Root length density used as index of BPP

¹Citations for different designs are: For Brookhaven National Laboratory (BNL) design, see Lewin *et al.* (1994) and Hendrey *et al.* (1999); for pure CO₂ injection design, see Miglietta *et al.* (2001a, 2001b) and Pepin & Körner (2002); for Screen Aided CO₂ Control design, see Leadley *et al.* (1997). ²Atmospheric CO₂ concentration (μmol mol⁻¹) that site strived to maintain and the time period (daylight hours only or full 24-h day, during growing season only or 365 d yr⁻¹) that CO₂ control was maintained. ³For precipitation data, station names and numbers from the US NOAA National Climate Data Center (NCDC) and UN World Meteorological Observatory (WMO) networks are given where off-site climate data was used. ⁴Latitudes and longitudes are: Finland 62°47' N, 30°56' E; Sweden 57°8' N, 14°30' E; Netherlands 51°59' N, 5°42' E; Switzerland 47°13' N, 7°3' E.




Fig. 3 Relationship between relative leaf CO₂ assimilation (A_{net}) and [CO₂] in the intercellular air spaces (C_i) composited across five woody species at three different free-air CO₂ enrichment (FACE) sites, using data from D. S. Ellsworth *et al.* (unpublished). A_{net} at all CO₂ levels is expressed relative to the CO₂ and light-saturated assimilation rate. (a) The overall relationship and linear extrapolation using $C_i/C_a = 0.70$ with two different model fits: (1) a hyperbolic fit and (2) fit from the Farquhar *et al.* (1980) model. The inset shows the two model fits in more detail. (b) Nonrectangular hyperbola function fit to all the data across species using measurements at 28°C for these curves. This temperature is used as it is close to the summertime mid-day temperature for FACE sites in North America.

oxygenation. However, a number of questions remain concerning the longer-term responses of A_{net} to increased [CO₂]: (1) what is the magnitude of the A_{net} response for field-grown plants; (2) is downregulation of photosynthetic metabolism significant; and (3) is the long-term A_{net} response sustained under plant resource limitations? In this section, we present some recent progress towards understanding the variability in CO₂ enhancement responses from studies of different plant species in FACE experiments.

The Farquhar *et al.* (1980) photosynthesis model provides a reference point to evaluate the response of A_{net} to elevated [CO₂]. Assuming that carboxylation rate is strictly CO₂-limited, one might expect that enhancements in A_{net} would be linearly proportional to the experimental enhancement in [CO₂] (i.e. the linear extrapolation in Fig. 3a). However, the observed short-term response of A_{net} to intercellular air space [CO₂] (C_i) (commonly called the $A_{net}-C_i$ response curve) is curvilinear, which indicates that strict CO₂-substrate limitation may not extend much higher in C_i than that commonly corresponding

with atmospheric [CO₂] (C_a). The two major components of the photosynthetic dark reactions in the Farquhar photosynthesis model (Farquhar *et al.*, 1980), carboxylation capacity under Rubisco-limited conditions (V_{c-max} prescribed by the Farquhar model) and RuBP regeneration resulting from electron transport (J_{max}), are assumed to virtually always scale with one another, including leaves grown under elevated CO₂ (Leuning, 1997; Medlyn *et al.*, 1999). Therefore, the family of $A_{net}-C_i$ response curves in all C₃ species should collapse to a single characteristic curve if normalized by leaf N-content or light- and CO₂-saturated maximum assimilation (A_{sat}) (Fig. 3b; Ollinger *et al.*, 2002; D. S. Ellsworth *et al.*, unpublished). We used two different curvature assumptions for the $A_{net}-C_i$ curve depicted in Fig. 3a: (1) an empirical rectangular hyperbola function (Hanson *et al.*, 1987; Ellsworth *et al.*, 1995) fit to data for multiple tree species in FACE experiments at 28°C; and (2) the theoretical model of Farquhar *et al.* (1980) fitted to data via the V_{c-max} and J_{max} parameters, also at 28°C. In both cases, for an enhancement in atmospheric [CO₂]




Fig. 4 Enhancement of CO_2 assimilation (A_{net}) for individual species from eight free-air CO_2 enrichment (FACE) sites in response to a 50–55% increase in growth $[\text{CO}_2]$. Cross-site measurements by D. S. Ellsworth *et al.* (unpublished) are grey bars, and measurements collected by specific site studies in the literature (see Table 1) are closed bars. Woody species are grouped on the left; herbaceous species on the right. Potential enhancements in A_{net} are shown as a ratio of A_{net} at elevated $[\text{CO}_2]$ divided by A_{net} at ambient $[\text{CO}_2]$, with the dashed line indicating no increase in A_{net} at elevated $[\text{CO}_2]$ as a reference point. Abbreviations for species: Liq, *Liquidambar*; Pin, *Pinus*; Pop, *Populus*; Bet, *Betula*; Amb, *Ambrosia*; Lar, *Larrea*; Lol, *Lolium*; Bro, *Bromus*; Koe, *Koeleria*; Agr, *Agropyron*; Tri, *Trifolium*; Sol, *Solidago*. Abbreviations for sites: TN, ORNL; NC, FACTS I; IT, PopFACE; WI, FACTS II; NV, NDFF; CH, ETH-Z; MN, BioCON; NZ, NZGraz.

from 365 to 565 $\mu\text{mol mol}^{-1}$, which is typical for many FACE sites, and assuming a $C_i : C_a$ ratio of 0.70, the response ratio E/A for A_{net} is either (69% of $A_{\text{sat}})/(49\% \text{ of } A_{\text{sat}}) = 1.41$ as predicted by (1), or (73% of $A_{\text{sat}})/(49\% \text{ of } A_{\text{sat}}) = 1.49$ as predicted by (2) (Fig. 3, inset). D. S. Ellsworth *et al.* (unpublished) observed a mean instantaneous response of 1.51 for photosynthesis in leaves with an instantaneous step change from 360 to 560 $\mu\text{mol mol}^{-1}$. Thus, theory and empirical measurements delineate the expected short-term enhancement of photosynthesis by elevated $[\text{CO}_2]$ as a basis against which the possibility of longer-term changes in photosynthetic enhancement can be evaluated.

A compilation of results for elevated $[\text{CO}_2]$ effects on A_{net} for 15 species from eight FACE sites indicates fairly substantial increases in A_{net} in response to a 50–55% increase in $[\text{CO}_2]$, particularly for woody plants (Fig. 4). Reviews of earlier studies under controlled-environment and OTC conditions, including several meta-analyses, also indicate enhancement of A_{net} at elevated $[\text{CO}_2]$ in herbaceous species and many woody plants (Drake & Leadley, 1991; Curtis, 1996; Medlyn *et al.*, 1999, 2001). Mean stimulation in A_{net} at the species level across all FACE literature studies was $26 \pm 5\%$ (range 0–60%; Fig. 4). It is important to bear in mind that this enhancement was typically measured under favorable environmental conditions during the growing season, and that diurnal and seasonal integrated enhancements in A_{net} (Ainsworth *et al.*, 2003; Naumburg *et al.*, 2003) may be different from those for physiological optimum conditions. Nonetheless, stimulation of A_{net} (based on species means, not averaged over the entire leaf area of the community) was variable in FACE experiments, ranging from essentially no stimulation in Minnesota prairie species (Lee *et al.*, 2001) to more substantial increases for pine and

sweetgum and large (45–60%) increases for *Populus* species on rich, irrigated soils (Fig. 4). Species in a Mojave Desert scrub (Naumburg *et al.*, 2003) and Wisconsin hardwood plantation (Noormets *et al.*, 2001) had intermediate stimulation values of 22% and 38%, respectively.

The A_{net} enhancement measured under naturally occurring conditions at FACE sites is smaller than that for controlled-environment and OTC studies (e.g. 53% in Saxe *et al.*, 1998, 51% in Medlyn *et al.*, 1999). However, $[\text{CO}_2]$ targets that are commonly used in FACE studies also differ from those in OTC studies. A coarse adjustment of the data presented in Medlyn *et al.* (1999) from model parameters suggests that a +22% response in A_{net} would be anticipated for about +55% elevated $[\text{CO}_2]$, which in turn indicates similarities in adjustments of the physiological responses of OTC and FACE plants (i.e. 22 and 26%, respectively) to long-term compared with short-term elevated $[\text{CO}_2]$ when considered to the same $[\text{CO}_2]$ target level. Note that these modeled A_{net} responses to elevated $[\text{CO}_2]$ are instructive for simple, coarse comparisons among different types of studies, but they cannot be used to indicate actual responses because the actual $[\text{CO}_2]$ attained is less than the target $[\text{CO}_2]$, as discussed above.

Because different sites may collect gas exchange data in slightly different ways, D. S. Ellsworth *et al.* (unpublished) collected A_{net} data across five woody taxa from three FACE sites in a standardized manner. When their data were compared with gas exchange measurements made by site investigators, A_{net} had a mean stimulation of $22 \pm 7\%$, which is similar to the average stimulation from previous literature measurements. In such a comparison, considerations such as measurements at different times of year and that stomatal conductance in particular is highly seasonally variable (Ellsworth, 2000; Nowak

et al., 2001) might explain some of the discrepancies between data sets. However, both sets of studies identified those species that were moderately and highly responsive to elevated $[\text{CO}_2]$, such as *Liquidambar*.

For a number of studies, the relative increase in A_{net} for leaves grown under ambient $[\text{CO}_2]$ v. leaves grown under elevated $[\text{CO}_2]$ was less than the instantaneous response to changes from 360 to 560 $\mu\text{mol mol}^{-1}$ $[\text{CO}_2]$ for leaves grown under ambient $[\text{CO}_2]$. The fact that the per cent increase in A_{net} was less than the expected per cent increase based on the Farquhar photosynthesis model suggests that biochemical adjustments in photosynthetic capacity (frequently termed 'down-regulation'; Sage, 1994) are important in governing plant production responses to elevated CO_2 . Such a reduction occurs as a result of changes in the operating C_i for gas exchange or down-regulation. Most evidence points to the latter phenomenon rather than the former (Medlyn *et al.*, 2001; von Caemmerer *et al.*, 2001). In fact, Huxman *et al.* (1998), von Caemmerer *et al.* (2001), Rogers & Ellsworth (2002), Ainsworth *et al.* (2003) and Bernacchi *et al.* (2003) all presented statistical evidence of downregulation in selected species in FACE experiments. However, in the majority of these studies, downregulation was observed under particular conditions such as low nutrient conditions (von Caemmerer *et al.*, 2001; Ainsworth *et al.*, 2003), in wet years when N and water are available but not in dry years (Huxman *et al.*, 1998; Naumburg *et al.*, 2003), and in older but not young leaves for evergreen species (Griffin *et al.*, 2000; Rogers & Ellsworth, 2002). By contrast, data for *Liquidambar* (Herrick & Thomas, 2001; Gunderson *et al.*, 2002), for *Populus tremuloides* (Takeuchi *et al.*, 2001) and for two *Populus* species (Bernacchi *et al.*, 2003) do not support an interpretation of photosynthetic downregulation in these species (Table 2). Therefore, the photosynthetic downregulation response appears to be both growth-form and environment specific. Nonetheless, even in experiments where downregulation is observed, A_{net} under elevated $[\text{CO}_2]$ is still stimulated in long-term experiments (Medlyn *et al.*, 1999), even after 7–10 yr of treatment (Ainsworth *et al.*, 2003; Crous & Ellsworth, 2004).

The maintenance of leaf enzyme content is critical to the sustainability of photosynthetic responses to elevated $[\text{CO}_2]$, and reductions in leaf N-content (Yin, 2002) or the rate of carboxylation per unit N are implicated when photosynthetic downregulation occurs (Medlyn *et al.*, 1999). We used the familiar relationship between A_{net} and leaf N (Reich *et al.*, 1998; Peterson *et al.*, 1999) to provide a framework for understanding the enhancement of A_{net} as a function of changes in leaf N-content, where leaf N-content serves as a proxy for photosynthetic protein content. Overall reductions in leaf N under elevated $[\text{CO}_2]$ would tend to reduce A_{net} , and reallocation of leaf N from photosynthetic to nonphotosynthetic leaf constituents would tend to reduce A_{net} and the ratio of $A_{\text{net}} : \text{N}$ in particular (Peterson *et al.*, 1999). However, the former would indicate a simple dilution




Fig. 5 (a) Relationship between A_{net} and leaf N content for plant species growing under ambient $[\text{CO}_2]$ at different free-air CO_2 enrichment (FACE) sites. Each point is a different species, and different symbol shapes are the different FACE sites shown in (b). Solid line is a simple linear regression for data except for the two inverted triangles enclosed by the oval; these represent two species that do not fit the overall relationship. Open symbols are herbaceous species and closed symbols are woody species. (b) Enhancement ratio of A_{net} at elevated $[\text{CO}_2]$ as a function of changes in leaf N content (leaf N content at elevated $[\text{CO}_2]$ divided by leaf N content at ambient $[\text{CO}_2]$) for species from six FACE sites. Solid line is a simple linear regression for data. Each point is a different species, and different symbol shapes are the different FACE sites. Open symbols are herbaceous species and closed symbols are woody species. FACE sites: open triangle, NZGraz; open square, BioCON; open circles, ETH-Z; closed circles, FACTS I; closed diamond, FACTS II; closed downward-pointing triangle, NDF; closed square, SAT; closed upward-pointing triangle, ORNL.

effect that might be a result of accumulation of simple carbohydrates in leaves under elevated $[\text{CO}_2]$, whereas the latter would indicate biochemical adjustments in leaves, as might occur owing to specific, hypothesized molecular mechanisms (Stitt & Krapp, 1999; Rogers & Ellsworth, 2002; Luomala *et al.*, 2003). A strong overall relationship between A_{net} at current $[\text{CO}_2]$ and leaf N-content occurred across four FACE sites (Fig. 5a; $P = 0.001$; $R^2 = 0.532$). However, the desert shrub species were outliers to this relationship because of N

Table 2 Summary of the most-recent published growing season results for light-saturated leaf net CO₂ assimilation (A_{net}) rate of C₃ plant species in free-air CO₂ enrichment (FACE) experiments

FACE site	Species	Exposure duration (yr)	Sampling date ¹	Measurement temperature (°C) ²	Mean (SE) ambient A _{net} ³	E/A for A _{net} ⁴	Response ratio N _{elev} : N _{amb}	Reference
NZGraz	<i>Trifolium subterraneum</i>	2	November 1999	20	20.1 (1.4)	1.08	0.912	von Caemmerer <i>et al.</i> (2001)
NZGraz	<i>Trifolium repens</i>	2	November 1999	20	20.1 (1.2)	1.06	0.8	von Caemmerer <i>et al.</i> (2001)
NZGraz	<i>Lolium perenne</i>	2	November 1999	20	15.6 (1.7)	1.40	1.018	von Caemmerer <i>et al.</i> (2001)
FACTS I	<i>Pinus taeda</i>	2.75	May 1999	28	5.3 (0.4)	1.25	0.99	Rogers & Ellsworth (2002)
FACTS I	<i>Liquidambar styraciflua</i>	3	September 1999	30 ± 0.5	13.2 (0.4)	1.58	1.17	Herrick & Thomas (2001)
BioCON	<i>Achillea millefolium</i>	1	August 1999	27.3	15.9 (0.7)	1.01	–	Lee <i>et al.</i> (2003)
BioCON	<i>Agropyron repens</i>	1	August 1999	27.3	9.0 (1.6)	1.03	1.14	Lee <i>et al.</i> (2003)
BioCON	<i>Bromus inermis</i>	1–2	July 1999	26.6	9.3 (1.9)	1.12	0.75	Lee <i>et al.</i> (2001)
BioCON	<i>Koeleria cristata</i>	1–2	July 1999	26.6	14.7 (1.6)	1.11	0.89	Lee <i>et al.</i> (2001)
FACTS II	<i>Populus tremuloides</i>	1.5	July 1999	Ambient ± 2	26	1.26	1.09	Takeuchi <i>et al.</i> (2001)
ORNL	<i>Liquidambar styraciflua</i>	2	July 2000	26 ± 2	12.4 (1.9)	1.53	1.1	Gunderson <i>et al.</i> (2002)
ETH-Z	<i>Lolium perenne</i>	9	May 2002	15 (air temp.)	16.8 (0.9)	1.45	–	Ainsworth <i>et al.</i> (2003)
NDFF	<i>Larrea tridentata</i>	4	April 2001	20	12.1 (1.1)	1.02	0.93	Naumburg <i>et al.</i> (2003)
NDFF	<i>Ambrosia dumosa</i>	4	April 2001	20	16.4 (4.2)	1.27	0.84	Naumburg <i>et al.</i> (2003)
SAT	<i>Larix decidua</i>	0.25	August 2001	21.1	10.1 (0.3)	1.46	0.95	Hättenschwiler <i>et al.</i> (2002)
PopFACE	<i>Populus alba</i>	3	July 2001	25–30	18.5 (2.5)	1.55	–	Bernacchi <i>et al.</i> (2003)
PopFACE	<i>Populus nigra</i>	3	July 2001	25–30	18.6 (2.7)	1.58	–	Bernacchi <i>et al.</i> (2003)
PopFACE	<i>Populus × euramericana</i>	3	July 2001	25–30	18.9 (1.4)	1.44	–	Bernacchi <i>et al.</i> (2003)

¹A mid- to late growing season sampling date was selected in cases where multiple measurements were made across a season. ²Measurement temperature is leaf temperature unless noted.

³Means (SE) of A_{net} for plants grown at ambient [CO₂]. ⁴The response of A_{net} to elevated [CO₂] treatment is defined as the ratio of leaf A_{net} in elevated [CO₂]-grown plants to A_{net} in ambient [CO₂]-grown plants.

stacking in thick leaves and lower photosynthetic nutrient-use efficiency. Of greater interest, however, is the relationship between the long-term $[\text{CO}_2]$ responses of A_{net} and leaf N-content (both area-based) across six FACE sites (Fig. 5b; $P < 0.001$; $R^2 = 0.568$). Species that showed the greatest reductions in leaf N-content at elevated $[\text{CO}_2]$ also tended to show reduced enhancement in A_{net} (Fig. 5b). This A_{net} -leaf N relationship would, of course, be expected, but it is indeed encouraging to see that data across multiple growth forms and sites support this expectation. Given that $A_{\text{net}}-C_i$ curves in diverse species collapse to a single relative response function that predicts a common enhancement of A_{net} (Fig. 3) and that variation in A_{sat} represents a range of leaf N contents, the correlation of the A_{net} response to elevated CO_2 with the leaf N response (Fig. 5) strongly suggests that resource availability mediates leaf N content and how it is maintained in ecosystems at elevated CO_2 , consistent with the original hypothesis of Strain & Bazzaz (1983).

Of additional importance is the clear dichotomy between herbaceous and woody species. Realizing that they are from different sites, most herbaceous species (except *Lolium*) show a reduction in leaf N content (mean reduction of 14%) and thus a modest mean enhancement in A_{net} of 12% at elevated $[\text{CO}_2]$, whereas woody species show a stronger enhancement in A_{net} of 38% but virtually no mean change in leaf N content over all species (Fig. 5b). Herbaceous species may indeed be more functionally plastic to a change in $[\text{CO}_2]$, as exemplified by a strong response in leaf N content, whereas woody species tend to exhibit a more conservative response.

V. Primary productivity

1. Datasets and statistical analyses

Measurements of production used in our datasets were from either destructive harvest of plant biomass or derived through allometric techniques; details on how data were collected and sources of the data are noted in the references in Table 1. We treated each measurement of above-ground, below-ground, or net (i.e. total above-ground plus below-ground) primary production for 1 yr at a site (abbreviated as APP, BPP and NPP, respectively) as an observation or replicate for our statistical analyses. Results from each of the two experiments at FACTS I and SCG were used as two independent studies. It is important to note that the $[\text{CO}_2]$ effect was not always significant for each year at each site, which raises the issue of how one treats data when the $[\text{CO}_2]$ effect is not significant. One approach is to set E/A equal to 1.0 when the $[\text{CO}_2]$ effect is not significant. However, this approach ignores the problems of high, naturally occurring variation and low replication that are typical for field experiments, both of which lead to an inherently low statistical power. Furthermore, this approach masks the real variation that occurs in natural systems and precludes any analyses that try to tease out

underlying causes of that variation, which is a goal of our review. Another approach is to use only those results that are significant. However, this approach yields a biased result that overestimates the true effect. A third approach, and the one that we use in this review, is to use all raw data without any alterations. The underlying assumption of this approach is that each data point is an estimate of the treatment effect. Although we have explicitly recognized limitations of the data sets (end of Section III above), each FACE experiment has been carefully designed to minimize potentially confounding effects. Thus, the raw data without any alterations represent the best available estimates of the $[\text{CO}_2]$ effect.

To statistically analyse the production data, we used a general linear model that included both discrete and continuous variables. Because we were interested in potential differences among vegetation types, vegetation type was the discrete, indicator variable in the regression model. Because we anticipated that production may vary with annual precipitation and that production may decrease through time, annual precipitation and years of CO_2 treatment were included in the regression model as continuous, predictor variables. Finally, we anticipated that the effects of precipitation or years of CO_2 treatment on production might differ among the vegetation types, so the interactions between vegetation type and each predictor variable were included in the model. Data were analysed with the PROC GLM procedure in SAS V8.02 (SAS, 2001) using the REGDIAG macro (Fernandez, 2003a).

Statistical output includes a typical ANOVA table and typical regression statistics (Table 3a). The ANOVA table indicates which variables in the linear model are significant, and interpretation of the significant model variables depends on if the variable is an indicator or predictor variable. A significant 'vegetation type' variable indicates a significant difference among vegetation types after linear relationships with the two predictor variables, annual precipitation and years of CO_2 treatment, are accounted for. Significant predictor variables indicate that a significant linear relationship occurred between production and the predictor variable across all vegetation types. A significant interaction between vegetation type and one of the predictor variables indicates that the linear relationship between production and the predictor variable differed among vegetation types. In addition to these statistical assessments, the P -value from the F -test of the overall linear model and the R^2 for the overall model are also reported.

We also ran 'reduced model' statistical analyses when the main effect of a predictor variable was not significant (Table 3b,c). For example, the main effect for the predictor variable 'years of CO_2 treatment' was not significant in the full model analysis of APP (Table 3a). Thus, this term and its interaction with vegetation type were dropped from the linear model, and data reanalysed with a reduced model where precipitation was the only predictor variable (Table 3b).

During examination of the results, we observed that this linear model over all data did not accurately characterize the

Table 3 Results from statistical analyses of above-ground (APP), below-ground (BPP) and net (NPP) primary production data from ecosystem free-air CO₂ enrichment (FACE) experiments

Variable	APP		BPP		NPP	
	df	P	df	P	df	P
(a) Full model ¹						
Vegetation type	2	0.238	1	0.464	1	0.012
Precipitation	1	0.011	1	0.405	1	0.060
Year CO ₂	1	0.366	1	0.258	1	0.347
Vegetation × precipitation	2	0.022	1	0.281	1	0.008
Vegetation × year CO ₂	2	0.428	1	0.336	1	0.016
Error	44		14		19	
Regression model P	0.003		0.069		0.001	
Regression R ²	0.414		0.582		0.675	
(b) Reduced model: annual precipitation only						
Vegetation type	2	0.169	2	0.457	1	0.248
Precipitation	1	0.003	1	0.792	1	0.047
Vegetation × precipitation	2	0.003	2	0.693	1	0.073
Error	47		16		21	
Regression model P	< 0.001		0.114		0.002	
Regression R ²	0.391		0.438		0.546	
(c) Reduced model: year CO ₂ only						
Vegetation type			2	0.710	1	0.972
Year CO ₂			1	0.833	1	0.359
Vegetation × year CO ₂			2	0.194	1	0.078
Error			16		21	
Regression model P			0.041		0.005	
Regression R ²			0.522		0.497	

¹Data were analysed with the general linear model procedure in SAS V8.02 (SAS, 2001) using the REGDIAG macro (Fernandez, 2003a) with the full model of vegetation type as a discrete (i.e. indicator or classification) variable and both annual precipitation and years of CO₂ treatment as continuous (i.e. predictor) variables (a), or with reduced model of vegetation type and only one of the continuous, predictor variables (b and c). *F*-tests were based on Type III sums of squares.

subset of data for APP of the forest sites. The linear relationships between APP and annual precipitation appeared to vary among the different forest sites, and the site-specific relationships differed from that across all forest sites. Thus, we also conducted 'reduced data set' analyses where we analysed APP data from only the forest sites in a separate general linear model (Table 4a). This statistical model was similar to that above except that FACE site was used as the discrete, indicator variable rather than vegetation type. As with the complete data set analyses, we also ran reduced model analyses that dropped nonsignificant variables from the linear model (Table 4b). For completeness, we also analysed APP data from only the grassland sites in a similar manner, but neither the site factor nor the site interaction terms were significant (results not shown).

Skewness, kurtosis, heteroscedasticity and normality of the data sets were examined for each statistical analysis. For all analyses except one, none of these ANOVA criteria were significant ($P \leq 0.05$). The only data set that failed these criteria was the NPP data. In 1994, NPP for the natural experiment at the SCG site was much lower than any other reported values, and this data point was highly influential on the statistical analyses

(as measured by its DFFITS statistic of -3.55) and appeared to be an outlier (as measured by its studentized residual of -3.78). We first tried several transformations of the data (ln, squared, etc.) to normalize the data set, but no transformation resulted in the data set passing the criteria. However, once this data point was removed from the NPP data set, skewness, kurtosis, heteroscedasticity and normality were no longer significant. Thus, we assumed that this data point was an outlier, and the statistical results given in this review exclude this data point. For completeness, we have shown the data point as an 'x' in Figs 6, 7, and 10.

2. Comparisons of ecosystems

As measured by either direct harvest or allometry-based techniques (Table 1), APP was almost always greater under elevated [CO₂] than under ambient [CO₂] (Fig. 6, top graph). Averaged over all 56 observations among 18 experiments at 16 FACE sites, APP under elevated [CO₂] was *c.* 19% greater than that under ambient [CO₂]. The greatest single-year increase in APP occurred in a desert ecosystem (E/A = 1.82, or 82% increase averaged over three shrub, two perennial grass, and




Fig. 6 Box plots of annual primary production, expressed as an enhancement ratio of production under elevated $[\text{CO}_2]$ to that under ambient $[\text{CO}_2]$, from 18 experiments at 16 free-air CO_2 enrichment (FACE) sites grouped by vegetation type. The thick line is the mean of the observed values, the thin line is the median value, the upper and lower lines of the box are the 75th and 25th percentiles, the upper and lower whisker lines are the 90th and 10th percentiles, and circles are potential outliers. Sample sizes are given in parentheses above the lower x-axis of each graph; the first number is the number of experiments with measurements and the second number is the total number of observations (an observation is 1 yr of production from an experiment, except for BioCON, where an average over 2 yr was the only result available). The '+' in below-ground primary production (BPP) of deserts indicates the actual observations (insufficient data were available to create a box plot), and 'x' in net primary production (NPP) of grasslands indicates a statistical outlier. Preplanned, least-squared comparisons among the forest, grassland,

four annual species; Smith *et al.*, 2000; Housman, 2002), followed by a forest plantation ($E/A = 1.71$ for leaf area index of poplar seedlings during their first year of growth; Gielen *et al.*, 2001). Pre-planned comparisons of ecosystems showed significant differences among desert, grassland, and forest ecosystems. On average, and when corrected for precipitation effects, increases in APP with elevated $[\text{CO}_2]$ were significantly greater in desert than in forest and grassland ecosystems ($P < 0.001$ for both comparisons) and greater in forest than in grassland ecosystems ($P = 0.055$), but the variation in E/A was also high for all ecosystems. Note that least-squared comparisons could not be made between the bog FACE sites and other ecosystems based on the general linear model (Table 3) because precipitation data for the bog ecosystems were not comparable to that of the other ecosystems. Using a simple one-way ANOVA of the data set and least significant difference comparison of means, mean APP of bogs was significantly less than that of deserts and forests ($P < 0.05$), but not different from that of grasslands (results not shown).

Not all sites measured BPP (Fig. 6, middle graph), and for those sites that did, the data have not been published for all years at all sites. It is very difficult to quantify BPP accurately (Böhm, 1979), and different techniques were used to estimate it among sites. Thus, the larger variation in E/A for BPP than for APP may be partly related to these measurement difficulties and differences. Forests generally had greater increases in BPP under elevated $[\text{CO}_2]$ than the other ecosystems (average E/A of 1.70 for forests vs 1.21, 1.11 and 0.93 for bog, grassland, and desert, respectively), but vegetation type was not significant in any of our statistical analyses (Table 3). The greatest increase in BPP (174%) was for a *Liquidambar* forest at the ORNL site during the fifth year of elevated $[\text{CO}_2]$ treatment (R. J. Norby, unpublished), whereas the greatest decrease in BPP (60%) was noted in an infertile, calcareous grassland during the first full year of treatment (Leadley *et al.*, 1999). The magnitude of this decrease may be inflated by random error, as subsequent years of results for the same site showed small (4–9%) increases in BPP at elevated $[\text{CO}_2]$. Nonetheless, decreased BPP was observed in another experiment at the same site (Niklaus *et al.*, 2001) as well as in an annual grassland (Shaw *et al.*, 2002).

Most ecosystems have greater NPP under elevated $[\text{CO}_2]$ (Fig. 6, bottom graph). Averaged over all 27 observations among 11 FACE sites that reported NPP, NPP was increased

and desert vegetation types were made based on the general linear model analysis in Table 3; note that the bog vegetation type could not be compared statistically with other vegetation types using this model because bogs did not have comparable precipitation data. Mean comparison within each graph are shown by the lower-case letters above each box; vegetation types within a graph with different letters were significantly different ($P < 0.05$). Note differences in ranges and scales for the y-axes among the production measurements. Dotted line across each graph shows ratio of response under elevated $[\text{CO}_2]$ to that under ambient $[\text{CO}_2]$ ($E/A = 1.0$ for reference). Sources for data are given in Table 1.

by almost 12% under elevated $[\text{CO}_2]$. Vegetation type was significant in the statistical analysis of NPP (Table 3a), and forests had a significantly greater increase in NPP with elevated $[\text{CO}_2]$ than did grasslands ($P < 0.05$). As with APP, bog ecosystems could not be compared with the other ecosystems using the general linear model. Using a simple one-way ANOVA of the data set and least significant difference comparison of means, mean E/A for NPP of bogs was significantly less than that of forests, but not different from that of grasslands. The maximum reported increase in NPP in any one year was approx. 38% in a forest ecosystem (R. J. Norby, unpublished). Decreased NPP under elevated $[\text{CO}_2]$ only occurred in grasslands, with three different experiments reporting $E/A < 1.0$ in one year each. The minimum E/A was 0.64 (i.e. a decrease in NPP by 36%) (Leadley *et al.*, 1999), and as discussed above, this point appeared to be a statistical outlier.

These results for the four different ecosystem types provide limited support for the early prediction that the response of productivity to elevated $[\text{CO}_2]$ would be greater in drier ecosystems (Strain & Bazzaz, 1983). The APP enhancement in elevated $[\text{CO}_2]$ was lowest in bogs and greatest in deserts, with forest and grasslands intermediate, which is consistent with the predictions of Strain & Bazzaz (1983). However, APP is only part of the carbon assimilated by plants, and results for NPP are contrary to the Strain & Bazzaz (1983) predictions. Although the results for BPP were not significantly different among ecosystems, E/A for BPP tended to decrease from mesic forest to xeric desert ecosystems. Phillips *et al.* (2002) speculated that higher water-use efficiency associated with elevated $[\text{CO}_2]$ and decreased stomatal conductance may allow sufficient water uptake with slightly smaller amounts of fine roots for plants growing under elevated $[\text{CO}_2]$ in deserts. Thus, plants in more water-limited environments may not need to invest as much carbon in their root systems to maintain water uptake and hence growth and productivity.

Interestingly, these overall increases in above-ground production for ecosystem FACE experiments are similar to those observed in crop FACE experiments (Kimball *et al.*, 2002). Excluding the results from the ETH-Z FACE site from Kimball *et al.*'s (2002) data set (ETH-Z results are also included in our data sets), crops with an ample supply of both water and N had an overall mean E/A for APP of 1.16, which is similar to the 1.19 value for all ecosystem studies. The elevated $[\text{CO}_2]$ effect on BPP for crops was substantially higher than that observed in ecosystems (average E/A for root biomass accumulation of grains and cotton and for potato tuber yield was 1.45, whereas BPP enhancement averaged over all available ecosystem data was 1.32).

3. Influence of water on enhancing production

A more rigorous test of the prediction that the relative effect of elevated $[\text{CO}_2]$ on growth increases as water availability

Table 4 Results from statistical analyses of above-ground production (APP) data for the reduced data set of only forest free-air CO_2 enrichment (FACE) sites

Variable	APP	
	df	P
(a) Full model ¹		
Site	3	0.418
Precipitation	1	0.042
Year CO_2	1	0.712
Site \times precipitation	3	0.702
Site \times year CO_2	3	0.024
Error	11	
Regression model P	< 0.001	
Regression R^2	0.903	
(b) Reduced Model: Annual precipitation only		
Site	3	0.178
Precipitation	1	0.002
Site \times precipitation	3	0.039
Error	15	
Regression model P	0.004	
Regression R^2	0.704	

¹Data were analysed with SAS V8.02 (SAS, 2001) using the REGDIAG macro (Fernandez, 2003a) with Site as an indicator (i.e. classification) variable and both annual precipitation and years of CO_2 treatment as predictor (i.e. continuous) variables in a general linear model (a) or for the reduced model that only had precipitation as the predictor variable (b). *F*-tests were based on Type III sums of squares.

decreases is to determine if $[\text{CO}_2]$ -induced changes in primary production is related to annual precipitation. Results from our general linear model analyses show that both APP and NPP are significantly correlated with annual precipitation, but these relationships differ among vegetation types (Table 3, Fig. 7). For forest ecosystems, the E/A for APP increases with increasing annual precipitation, but the relationship differs among FACE sites (Table 4; Fig. 7, top left panel). The E/A for APP also increases with increased precipitation for deserts, but it decreases for grasslands. The decreasing relationship for grasslands is consistent across all sites, with one regression fitting the data for all grassland sites. For NPP, E/A increases with increasing precipitation for both forests and grasslands (Table 3), but the slope of the relationship in forests is greater than that in grasslands. For all FACE sites, E/A for BPP is not significantly related to annual precipitation (Table 3). Thus, the only data set from ecosystem FACE experiments that is consistent with the prediction that the $[\text{CO}_2]$ enhancement of productivity will be greater in dry years is APP of grasslands; all other data sets either show the opposing trend or are not significantly related to precipitation. Interestingly, crops also show increased above-ground production under low water (but ample N) growth conditions, with E/A increasing from 1.16 for ample water conditions to 1.20 for low water conditions (Kimball *et al.*, 2002).




Fig. 7 Relationship between enhancement ratio of primary production and annual precipitation for forest (left three graphs), grassland (central three graphs), and desert (right two graphs) free-air CO_2 enrichment (FACE) sites. Top three graphs are above-ground, middle three are below-ground, and bottom two are net primary production; note differences in range and scaling for the y-axis among the production measurements. Each symbol represents the ratio of response under elevated $[\text{CO}_2]$ to that under ambient $[\text{CO}_2]$ (E/A) of production for 1 yr along with the corresponding precipitation for that year. Within each vegetation type, different symbol shapes represent different FACE experiments; the 'x' in the net primary production (NPP) of grasslands graph is a statistical outlier that was excluded from the statistical analyses (see text). Solid lines represent significant linear relationships between production and precipitation for all data within a graph, except for the upper left (above-ground primary production (APP) of forest) graph; for this graph, individual lines are regressions for four forest FACE experiments with > 3 yr of measurements (significant site \times precipitation term in the statistical analysis of the above-ground production data for the forest vegetation type; see Table 4b). The P and R^2 values for the linear regressions are shown in Tables 3 and 4. For data sets that did not have a significant linear relationship, the mean value of all observations is indicated by a dashed line. A dotted line across each graph shows E/A = 1.0 for reference. Sources for data are given in Table 1.

Results for APP data from grassland OTC experiments agree well with the ecosystem FACE data (Fig. 8, top graph). The solid lines in Fig. 8 are the same as those shown in the top panels of Fig. 7 and are color coded for the different vegetation types. The open green symbols are data from a short-grass (squares, Morgan *et al.*, 2003) and a tall-grass prairie (hexagons; Owensby *et al.*, 1999), and except for the driest year of OTC data, the regression line between APP E/A and annual precipitation for the OTC data is not significantly different from that for the grassland FACE data. Interestingly, the driest year of OTC data falls within the 95% confidence intervals around the regression line for the desert FACE site.

Comparisons between OTC and FACE results for forests are more difficult to make because the relationship between E/A for APP and precipitation is site-specific for the FACE sites. When annual precipitation is near 1500 mm, OTC data fall

well above the regression lines for the FACE data. However, these OTC data are for rapidly growing seedlings, whereas the FACE data are for older trees nearing or past canopy closure. In addition, the FACE site with the steepest slope consists of rapidly growing seedlings (FACTS II). Thus, the greater E/A from the OTC experiments may simply be related to differences in tree age and their accompanying developmental stage (Norby *et al.*, 1999).

Taken together, these results do not provide strong support for predictions that the response of productivity to elevated $[\text{CO}_2]$ would be greater in drier ecosystems or in dry years within an ecosystem. The difference in the APP-annual precipitation relationship between grassland and desert ecosystems is especially surprising because one would expect that these primarily water-limited ecosystems would have similar responses to elevated $[\text{CO}_2]$. Despite this apparent




Fig. 8 Comparison of above-ground production enhancements as related to annual precipitation between free-air CO_2 enrichment (FACE, solid lines) and open-top chamber (OTC, open symbols) experiments. Results for FACE experiments are the same regression lines shown in Fig. 7 except color-coded here (dark red, deserts; dark green, grasslands; dark blue, forests). Results for OTC experiments are from tall-grass prairie (light-green hexagons), short-grass-prairie (light-green squares), *Liriodendron tulipifera* (light-blue triangles) and *Quercus alba* (light-blue inverted triangles). Black dashed line is a conceptual model to explain the relationship between the ratio of response under elevated $[\text{CO}_2]$ to that under ambient $[\text{CO}_2]$ (E/A) for above-ground primary production (APP) and annual precipitation in ecosystems that are primarily water-limited.

discrepancy, we suspect that a more complex conceptual model fits the APP results for water-limited systems. This conceptual model, which is represented by the black dashed line in Fig. 8, top graph, has two components. First is the benefits of increasing water-use efficiency (WUE) from elevated $[\text{CO}_2]$ that underlies the predictions from Strain & Bazzaz (1983) – as exhibited by grasslands, the increasing benefits of increased WUE under elevated $[\text{CO}_2]$ results in a greater enhancement of APP as annual precipitation decreases. However, just as high WUE is not sufficient for survival of desert plants, the benefits of increased WUE from elevated $[\text{CO}_2]$ has only limited effects on APP as precipitation continues to decrease. As demonstrated by desert vegetation, the response of APP to elevated $[\text{CO}_2]$ is constrained at very low precipitation by the need for plants to cope with severe drought. Thus, the enhancements of APP by elevated $[\text{CO}_2]$ peaks at some intermediate precipitation (the empirical results from the FACE and OTC experiments suggest that the maximum E/A for APP occurs between 300 mm and 500 mm annual precipitation).

Results for forest ecosystems clearly do not fit predictions that enhancements of productivity increase with decreased water availability. The relationship of increased E/A with increased annual precipitation (Fig. 7) was highly significant and well correlated for both APP (Table 4b) and NPP (Table 3a). Although differences among sites in steepness of the slope for APP appear to be related to stand age, as discussed above, we still need to account for why E/A increases with increased precipitation, contrary to the early conceptual model of Strain & Bazzaz (1983). One explanation might be

that annual precipitation is not the best index of water availability for forest production. For example, even though the growth of large trees in wet years is greater than that in dry years, seasonal timing of rainfall has a large impact on their growth (Hanson *et al.*, 2001). Thus, if short-term drought at critical growth periods is accompanied by wet periods at other, noncritical times during the year, then the short-term drought will be masked in the annual precipitation measurements. A second explanation may be that nutrient availability increases with increased precipitation, which then drives the increased productivity response to elevated $[\text{CO}_2]$. However, recent studies of the N cycle at three forest FACE sites did not find any significant effects of elevated $[\text{CO}_2]$ on microbial N cycling (Zak *et al.*, 2003). Another possibility would be that some other climatic factor, such as growing season temperature or length of the growing season, covaries with annual precipitation. For example, photosynthetic enhancement by elevated $[\text{CO}_2]$ increased with increased leaf temperature for loblolly pine at FACTS I (Myers *et al.*, 1999).

4. Influence of nitrogen on enhancing production

Although fewer data are available, an enhanced $[\text{CO}_2]$ -effect on production typically occurs when ecosystem N is increased (Fig. 9), a response that is consistent with earlier predictions. The results, shown as wide, dark-colored bars in Fig. 9, are from three grassland, one forest, and one desert FACE site where soil N varied either by adding additional N (BioCON, ETH-Z, FACTS I Prototype, MEGARICH) or by natural variation among different microsites (NDFF). Note that for




Fig. 9 Effects of increased nitrogen availability on the enhancement of primary production by elevated $[\text{CO}_2]$. Wide, dark-colored bars are results from ecosystem free-air CO_2 enrichment (FACE) experiments whereas narrow, light-colored bars are from meta-analysis of controlled-environment and open-top chamber (OTC) experiments. The ratio of response under elevated $[\text{CO}_2]$ to that under ambient $[\text{CO}_2]$ (E/A) for low nitrogen (N) availability (closed bars) are production under elevated $[\text{CO}_2]$ and low N availability divided by production under ambient $[\text{CO}_2]$ and low N. The E/A ratio for high N availability (hatched bars) are production under elevated CO_2 and high N availability divided by production under ambient $[\text{CO}_2]$ and low N. Except for NDFF, results for ecosystem FACE experiments are means and standard errors, averaged over all years of observation at a site; only 1 yr of data from the NDFF was available. For the meta-analysis studies, results are recalculated using all controlled-environment and OTC experiments in the databases of the meta-analyses and are means and standard errors. A dotted line across each graph shows $\text{E/A} = 1.0$ for reference. Sources for ecosystem FACE data are given in Table 1.

the three sites where N was added, the N additions were within generally accepted management practices. The low N values in Fig. 9 represent the ratio of production under the combined treatments of elevated $[\text{CO}_2]$ and low N to that under the combined treatments of ambient $[\text{CO}_2]$ and low N, i.e. the same N treatment but different $[\text{CO}_2]$ treatments. To directly determine if increased N availability further increases the $[\text{CO}_2]$ effect, we need to use the same reference point for production under high N (i.e. we need to use the same denominator – combined treatments of ambient $[\text{CO}_2]$ and low N – but the numerator is now the combined treatments of elevated $[\text{CO}_2]$ and high N). Thus, if the $[\text{CO}_2]$ effect increases when additional N is available, then the height of the bars should increase from low to high N availability, which occurs without exception for the ecosystem FACE experiments.

To compare these ecosystem FACE results with controlled-environment and OTC experiments, we recalculated similar E/A ratios for low and high N studies using the databases from meta-analyses of trees (Curtis & Wang, 1998) and grasses (Wand *et al.*, 1999). Results from the controlled-environment and OTC studies show a similar pattern to the ecosystem FACE experiments, although the enhancement of NPP when N is more readily available is much greater for the controlled-environment and OTC studies than for the ecosystem FACE experiments. This much greater enhancement of E/A for NPP

in controlled-environment experiments is likely caused by the optimal growing conditions that plants experience during the experiments, the short periods during which the experiments are conducted and the fact that most experiments use seedlings that are rapidly growing. Results from other OTC experiments that are not included in these meta-analyses (Norby *et al.*, 1999; Joel *et al.*, 2001) show increased enhancements of productivity when nutrients are more available that are more similar to those of FACE experiments. For example, the E/A values for APP under ambient N were 1.32 and 1.07 for sandstone and serpentine communities, respectively, and the enhancement ratios, expressed as APP under elevated CO_2 and high N vs APP under ambient CO_2 and ambient N, increased to 2.22 for sandstone and 1.55 for serpentine communities (Joel *et al.*, 2001). Further support for this greater enhancement of the $[\text{CO}_2]$ effect comes from crops: the E/A for APP of rice grown under high N was greater than that under ample N (Kimball *et al.*, 2002), although the E/A for BPP was less under high N.

Related to the prediction that increased N will increase the $[\text{CO}_2]$ effect is that ecosystem responses to a step-change in $[\text{CO}_2]$ are transient because ecosystems quickly develop N limitations (Luo & Reynolds, 1999), which in turn will decrease E/A over time. For the 10 FACE experiments that have at least 3 yr of results, neither the E/A for APP nor that for BPP significantly changed over time when precipitation




Fig. 10 Relationship between primary production and the number of years of CO_2 treatment for free-air CO_2 enrichment (FACE) experiments that have reported three or more years of above-ground production results. Top three graphs are above-ground, middle three are below-ground and bottom two are net primary production; left three graphs are for forest, center three for grassland, and right two are desert FACE sites. Note differences in range and scaling among the production measurements. Each point is production for the first, second, third, etc., year of CO_2 treatment. Within each vegetation type, different symbol shapes represent different FACE experiments; the 'X' in the net primary production (NPP) of grasslands graph is a statistical outlier that was excluded from the statistical analysis (see text). Solid lines represent significant linear relationships between production and years of CO_2 treatment for all data within a graph; the P and R^2 values for the linear regression are shown in Table 3. For data sets that did not have a significant linear relationship between production and years of CO_2 treatment, the mean value of all observations is indicated by a dashed line. A dotted line across each graph shows ratio of response under elevated $[\text{CO}_2]$ to that under ambient $[\text{CO}_2]$ ($E/A = 1.0$ for reference). Sources for data are given in Table 1.

effects were also accounted for (Table 3a; Fig. 10 top and middle panels). For the E/A of NPP, the vegetation type \times years of CO_2 interaction term was significant (Table 3a), indicating that the relationship between the NPP enhancement and years of CO_2 treatment for forests sites differed from that for grassland sites. For forests, the E/A for NPP significantly increased with time, whereas that for grasslands decreases (Fig. 10 bottom panels). Although the $[\text{CO}_2]$ effect on production does not appear to be dampened over time, except for NPP of grasslands, these results must be interpreted cautiously; results from models suggest that the decrease in E/A with time likely occurs on a time-scale of > 10 yr (Luo & Reynolds, 1999). Thus, it is critically important that the current FACE experiments be continued.

VI. Response of plant functional types


Free-air CO_2 enrichment studies, most of which use intact vegetation that is not highly manipulated experimentally, are not well suited for explicitly examining community diversity responses to elevated $[\text{CO}_2]$, especially given the long life-span of perennial dominants relative to the length of time that the experiments have been occurring. However, an examination of the relative responses of different plant functional groups can serve as a proxy for how changing $[\text{CO}_2]$ may influence

the relative abundance of these functional types over longer periods.

The data set to determine if the effects of elevated $[\text{CO}_2]$ differed among plant functional types consists of measurements of APP for individual species at different FACE experiments. In most cases, data are reported for an individual species, but we also used data reported for a group of species. When the data set for a functional type included a single mean reported for a group of species, we represented that result as a single data point in the data set (as opposed to replicating the mean a number of times in our data set to match the number of species that went into that mean). To indicate that a particular data set had data that included a single mean for a group of species, ' $>$ ' is placed before the number of species in Fig. 11, i.e. the sample size for the number of species is actually greater than the number of means in the data set. For many species, two or more years of observations are reported, and a replicate is considered to be an individual year of results for an individual species (or a group if that was the only reported data) for an individual experiment.

Statistical analysis to compare E/A for APP among plant functional types was complicated by three factors. First, not all sites have representatives of all functional types. Second, no one species was common among all sites. Third, measurements for most species were not made or reported in all years of

Fig. 11 Enhancement ratios of above-ground production for different plant functional types. Results from the ecosystem experiments in this review are shown as box plots where the thick line is the mean of the observed values, the thin line is the median value, the upper and lower lines of the box are the 75th and 25th percentiles, the upper and lower whisker lines are the 90th and 10th percentiles, and circles are potential outliers. Sample sizes for the box plots are given in parentheses after the plant functional type name; the first number is the number of free-air CO₂ enrichment (FACE) experiments with measurements, the second number is number of species with observations (note that '>' indicates that results used from the literature were averaged over more than one species), and the total number of observations (an observation is 1 yr of production for a species at a site). Preplanned, least-squared comparisons among the plant functional types for the ecosystem FACE experiments were not significant. Corresponding results for crop plants from FACE experiments as well as data from controlled-environment and open-top chamber (OTC) studies are shown as closed symbols. Dotted line across graph shows ratio of response under elevated [CO₂] to that under ambient [CO₂] (E/A) = 1.0 for reference.



the experiment. Thus, a simple repeated measures factorial ANOVA could not be used to examine the data because of the large number of missing cells in the data set. To overcome these issues, we first constructed a dummy variable called 'SiteType' that combined the specific site information with the plant functional type information (e.g. ShrubORNL would be results for shrub species at ORNL). Next, APP was ln-transformed so that the data set met skewness, kurtosis, heteroscedasticity, and normality criteria. Then, SiteTypes were compared using a Qualitative X Qualitative Repeated Measures ANOVA (Fernandez, 2003b) where SiteType and Species were the two qualitative factors and Year was the repeated measures factor; this analysis utilized PROC MIXED in SAS. Because the SiteType main effect was significant in this ANOVA (Table 5a), we then conducted a series of planned contrasts to examine if plant functional types differed from each other, where all sites with one functional type were compared with those for another functional type (Table 5b). One nonlegume, *Solidago canadensis* at ORNL, had E/A > 25 for each of the two years of measurements (Belote *et al.*, 2003), which results in the relatively high mean and 90th percentile for the nonlegumes. Although we emphasize that the ln-transformed data met all four assumptions of ANOVA's, we also analysed the data set without these two data points. Again, data had to be ln-transformed to meet the ANOVA criteria, but the SiteType main effect was not significant in this ANOVA ($P = 0.206$). Thus, we conclude that the data set

with the two *Solidago* data points does not bias the statistical analyses.

Because legumes fix N and hence potentially avoid severe N limitations, and because increased A_{net} under elevated [CO₂] could alleviate some of the carbon limitations for N-fixation, legumes have been predicted to have greater responses to elevated [CO₂] than other plant functional types (Tissue *et al.*, 1997; Grünzweig & Körner, 2001). Although the effects of elevated [CO₂] on APP of legumes tended to be greater than those reported for some other plant functional types (Fig. 11), the planned contrasts between legumes and all other functional types across all FACE sites was not significant (Table 5b). The four FACE experiments that had legumes also had nonleguminous herbs, and although the E/A ratio of the legumes was numerically greater than that of the nonlegumes for three of the experiments, the difference between legumes and nonlegumes was not significant for any one experiment (Lüscher *et al.*, 1998; Leadley *et al.*, 1999; Niklaus *et al.*, 2001; Reich *et al.*, 2001b). Thus, although legumes tended to have the expected greater response to elevated [CO₂], the variation in responses among sites (Fig. 11) as well as among species within each site suggest that the N-fixation trait is not sufficient to predict how leguminous and nonleguminous herbs will respond to elevated [CO₂].

Other comparisons of interest are those between C₃ and C₄ plants and between woody and nonwoody plants, where C₃ (and nonwoody) plants have been predicted to have a greater

Table 5 Results from statistical analyses of ln-transformed above-ground production (APP) for different plant functional types from ecosystem free-air CO₂ enrichment (FACE) experiments

Variable	Numerator df	Denominator df	<i>P</i>
(a) ANOVA ¹			
SiteType	32	36	< 0.001
Species	8	35	0.303
Year	4	80	0.696
(b) Planned contrasts			
C ₃ vs C ₄	1	37	0.770
C ₃ vs Legume	1	41	0.738
C ₃ vs Nonlegume	1	39	0.352
C ₃ vs Shrub	1	41	0.282
C ₃ vs Tree	1	34	0.929
C ₄ vs Legume	1	38	0.601
C ₄ vs Nonlegume	1	36	0.341
C ₄ vs Shrub	1	38	0.288
C ₄ vs Tree	1	34	0.824
Legume vs Nonlegume	1	41	0.639
Legume vs Shrub	1	42	0.510
Legume vs Tree	1	37	0.685
Nonlegume vs Shrub	1	40	0.786
Nonlegume vs Tree	1	34	0.324
Shrub vs Tree	1	36	0.262

¹Data were analysed with SAS V8.02 (SAS, 2001) using a 'QI × QI repeated' routine (Fernandez, 2003b). *F*-tests were based on Type III sums of squares. Numerator and denominator degrees of freedom are shown for each *F*-test.

response than C₄ (and woody) plants (Strain & Bazzaz, 1983). The C₃ grass functional type had a slightly greater mean response to elevated [CO₂] than C₄ grasses (Fig. 11), but this difference, as well as those between C₄'s and all other functional types, was not significant (Table 5B). Similarly, comparisons between woody functional groups (trees and shrubs) and the other functional groups were also not significant. Taken together, these results for different functional types suggest 'that current trait-based functional classifications might be useful, but not sufficient, for understanding plant and ecosystem responses to elevated CO₂' (Reich *et al.*, 2001b) and that environmental factors interact with plant functional type to influence plant responses to elevated [CO₂].

Results from controlled-environment and OTC experiments were near the mean or median of data from the ecosystem FACE experiments (Fig. 11). The controlled-environment and OTC data are from meta-analyses of woody plants (primarily trees; Fig. 11, closed triangles) or C₃ and C₄ grasses (closed, inverted triangles), from a shrub-oak OTC experiment (closed diamonds), or was compiled from various literature (closed hexagons). However, results for crops when grouped by functional types (closed squares) did not always match those for plants from ecosystem studies (Fig. 11). We compared data for crops, which are from Kimball *et al.* (2002), with

ecosystems as follows: cotton and grapes (woody perennials) are compared with shrubs; potato (broadleaf forb) with nonlegumes; wheat and rice (C₃ grains) with C₃ grasses; and sorghum (C₄ grain) with C₄ grasses. Mean E/A for crop functional types were numerically less than those for ecosystem functional types, but the data overlapped extensively. However, crops are not bred for APP, but for their agricultural yield, and E/A of agricultural yields typically are greater than those for APP, especially under water or N limitation (Kimball *et al.*, 2002).

Several lines of evidence suggest that elevated [CO₂] may preferentially increase the abundance of another group of plants – invasive species (Dukes & Mooney, 1999; Weltzin *et al.*, 2003). For example, results from a controlled-environment experiment with six common invasive species found that these plants were more responsive to [CO₂] increases that have occurred in the past century than to the rise in [CO₂] anticipated in the next century, suggesting that changes in [CO₂] have already played a stimulatory role in plant invasions (Ziska, 2003). The most responsive species to elevated [CO₂] in the desert was an invasive C₃ annual grass (Smith *et al.*, 2000), while production of another invasive species, a C₃ woody vine, increased threefold in a forested ecosystem at ORNL (Belote *et al.*, 2003). However, not all invasive species respond favorably to elevated [CO₂]: an invasive C₄ annual grass showed reduced production under elevated [CO₂] in the same ORNL experiment. However, this reduced production for the C₄ annual is not unexpected when compared with the responses of other C₄ plants (Fig. 11). Thus, the available results provide support for the prediction that elevated [CO₂] favors at least C₃ invasive species. Furthermore, when invasive species have a potentially disruptive role through a stimulation of anomalous change in the ecosystem, they could have a major impact on subsequent community structure and diversity. For example, *Bromus* spp. are known to stimulate a fire cycle (Sage, 1996) and to alter N-cycling (Evans *et al.*, 2001) in arid and semiarid ecosystems in the western USA. The differential stimulation of growth and seed production in *Bromus* by elevated [CO₂] could therefore result in a marked change in community structure and function of the ecosystems they are currently invading (Smith *et al.*, 2000).

More diverse plant communities are predicted to be more responsive to elevated [CO₂] (Bolker *et al.*, 1995). As demonstrated by the results above, along with others (Hooper & Vitousek, 1997; Tilman *et al.*, 1997), all plants are not equal in their responses to environmental perturbations, even those within a functional group. Thus, the prediction that communities with greater biodiversity will have a greater [CO₂] response may or may not be due to some inherent characteristic of biodiversity. For example, more diverse communities are more likely to have more species that respond more strongly to elevated [CO₂], which can be interpreted as a 'sampling' effect (Loreau, 2000; Niklaus *et al.*, 2001). Two FACE sites have explicitly examined the biodiversity hypothesis:

BioCON and SCG. Both studies showed significantly increased production for more diverse species assemblages. The elevated $[\text{CO}_2]$ effect changed from +7% to +22% as the number of species increased from 1 to 16 at BioCON (Reich *et al.*, 2001a). For the biodiversity experiment at the Swiss calcareous grassland, the $[\text{CO}_2]$ effect varied through time, but E/A ratios of total community biomass at final harvest were 0.80, 1.08 and 1.41 for communities that initially had 5, 12 or 31 species (Niklaus *et al.*, 2001). Some functional group effects were evident (Craine *et al.*, 2003), but in both experiments, the effects of biodiversity on $[\text{CO}_2]$ responses were largely caused by a subset of the species that were used in the experiments, which is consistent with the 'sampling effect' concept. Interestingly, biodiversity of a California grassland that is dominated by annual grasses decreased with elevated $[\text{CO}_2]$, primarily because of decreased forb diversity (Zavaleta *et al.*, 2003). Clearly, the influence of biodiversity, plant functional type composition, and species identity all play a role in ecosystem-level responses to elevated $[\text{CO}_2]$.

VII. Conclusions

Results from FACE sites (i.e. for plants growing in a competitive matrix under natural conditions) have largely substantiated predictions of how elevated $[\text{CO}_2]$ affects leaf CO_2 assimilation. As expected, almost all the C_3 species examined at different FACE sites have increased photosynthesis under elevated $[\text{CO}_2]$. Also as expected, the increase in A_{net} with elevated $[\text{CO}_2]$ when averaged over all species (26%) was less than the increase in $[\text{CO}_2]$ (50–55%), and the increase in ecosystem primary production (19% for APP, 32% for BPP, and 12% for NPP) was less than both. The enhancement of A_{net} for field-grown plants was roughly similar to that for plants in controlled-environment and OTC studies after adjusting for differences in experimental $[\text{CO}_2]$. Finally, downregulation of photosynthesis occurred in a number of FACE experiments, but not in all species and only under certain conditions for other species.

Although the effects of elevated $[\text{CO}_2]$ on A_{net} varied among species, two relationships were consistent. First, the enhancement of photosynthesis by elevated $[\text{CO}_2]$ was positively related to the relative change in leaf N content: smaller increases in A_{net} under elevated $[\text{CO}_2]$ occurred for plants that had reduced leaf N under elevated $[\text{CO}_2]$, whereas plants that showed no or small increases in leaf N had larger increases in A_{net} . Second, herbaceous species consistently had reduced leaf N-content under elevated $[\text{CO}_2]$, and hence smaller enhancements of A_{net} , than woody species. Thus, herbaceous species may be more functionally plastic than woody species under elevated $[\text{CO}_2]$. This assimilation–leaf N relationship also implies that the effects of elevated $[\text{CO}_2]$ on A_{net} is responsive to fertility or increased N deposition, and downregulation of photosynthesis may occur more prevalently under N-limited conditions, as earlier predicted. Inter-

estingly, the ecosystems that had the greatest enhancements of A_{net} (i.e. woody-dominated ecosystems such as forests and deserts) had the greatest enhancements in APP, suggesting that A_{net} may scale to the ecosystem level, although the form and value of the scaling factor is not currently known. Nonetheless, results for BPP and NPP suggest that the scaling factor must also include an allocation component, as responses of BPP to elevated $[\text{CO}_2]$ differ between forests and deserts.

Results for predictions of the effects of elevated $[\text{CO}_2]$ on primary production are more mixed, but are generally less than a 20% increase in NPP based on the β -factor approach. The results from FACE sites suggest that a single β is not feasible, at least for global predictive purposes, given differences among ecosystems types and differences in plant responses to elevated $[\text{CO}_2]$ in combination with other environmental parameters such as water and N availability. Early predictions that are based on resource limitations generally are valid for N availability, but less so for water availability. The significantly greater enhancement of APP for deserts compared with other ecosystems is consistent with predictions that enhancement of productivity would be greater in drier ecosystems, but the result that forests have significantly greater enhancements of APP and NPP than grasslands is not. Furthermore, only APP for grasslands had greater productivity enhancements in drier years – relationships between the enhancement of APP and annual precipitation for forests and deserts and those between BPP or NPP and annual precipitation were either not significant or increasing. By contrast, the predicted increase in productivity enhancement with increased N availability is well supported by the FACE results as well as by controlled-environment and OTC studies. Although the water availability effects are counter to initial predictions of Strain & Bazzaz (1983) made 20 yr ago, their initial conceptual model based on the importance of increased WUE under elevated $[\text{CO}_2]$ coupled with the concept of an over-riding importance of drought survival mechanisms at very low precipitation is consistent with an empirical fit to data from water-limited ecosystems (Fig. 8).

Results for plant functional type also do not generally fit predictions that certain groups of plants such as legumes and C_3 plants will have greater enhancements of productivity than woody and C_4 plants. Two factors are important in this assessment of plant functional types. First, predictions based on the $[\text{CO}_2]$ enhancement of productivity from single, isolated plants are not good predictors of how well those same plants do under interspecific competition (Poorter & Navas, 2003). Second, the identity of individual species is important, i.e. not all species within a functional type are necessarily similar. The presence or absence of individual species and functional groups can significantly influence responses of native ecosystems to elevated $[\text{CO}_2]$ and their feedbacks with other global change factors (Reich *et al.*, 2001a; Zavaleta *et al.*, 2003), and we anticipate greater attention to this topic in current and future experiments.

We have focused almost exclusively on plant processes in this review, primarily because the majority of results from FACE sites that have been reported to date concern plant processes and hence they form the most comprehensive data set. Fortunately, results from other trophic levels are beginning to emerge. For example, increases in soil microbial biomass (Schortemeyer *et al.*, 1996) and changes in N cycling (Billings *et al.*, 2002) are evident in some ecosystems, as postulated by Zak *et al.* (1993), although changes in soil microbial biomass or composition do not always change with elevated [CO₂] (Allen *et al.*, 2000; Zak *et al.*, 2000, 2003; Niklaus *et al.*, 2003). Changes in leaf quality with elevated [CO₂] have also been noted, although these changes had either no or only subtle effects on insect herbivores for aspen (Percy *et al.*, 2002; Kopper & Lindroth, 2003). Effects on plant fungal pathogens are also variable, with no effects noted for aspen (Percy *et al.*, 2002) but increased pathogen loads on C₃ grasses under elevated [CO₂] (Mitchell *et al.*, 2003). These variable, and sometimes surprising, results further emphasize Körner's (2000) conclusion that only a fully coupled ecosystem approach will yield meaningful information on how the biosphere may respond to global change. We contend that integrated studies at FACE sites are currently our best method to achieve this approach.

This review also has focused primarily on an elevated [CO₂] effect and largely ignored other environmental factors that are likely to change as [CO₂] continues to increase. With the exception of N availability (Fig. 9), multifactor FACE experiments are rare. Realistic, multifactor experiments are difficult and costly to achieve in some ecosystems such as forests and deserts, but the importance of factor interaction is evident from the studies in the annual grassland community at JRGCE (Shaw *et al.*, 2002). Although elevated [CO₂] alone increased NPP by *c.* 8%, the interactive effects of elevated [CO₂] with temperature, N, and precipitation on NPP were less than those of ambient [CO₂] with those factors. These results clearly point to the need for multifactor experiments, and they also strengthen the argument that both resource availability and species composition are important. Thus, an integrative approach, such as that proposed by Field *et al.* (1992), is needed to predict the impacts of a suite of global changes on ecosystem attributes.

Clearly, results from the FACE sites that are summarized in this review have greatly advanced our knowledge of plant and ecosystem responses to elevated [CO₂] for specific ecosystems as well as across multiple ecosystems. But the utility of these experiments goes well beyond simply predicting global change effects. These FACE experiments are often designed to investigate fundamental mechanisms that drive ecosystem structure and function, issues that are at the heart of ecology. Thus, the importance of these FACE sites is not only how well they can predict the impacts of elevated [CO₂], but also how well they test ecological concepts.

Acknowledgements

We thank Lesley DeFalco, Evan DeLucia, Bert Drake, Jeff Dukes, Chris Field, Dave Housman, Dave Karnosky, Mark Kubiske, Dave Moore, Jack Morgan, Elke Naumburg, Paul Newton, Rich Norby, Don Phillips, Jake Weltzin and Stephen Zitzer for sharing some of their data that they have incorporated into manuscripts for publication. We also greatly appreciate the statistical advice of George Fernandez and the constructive comments on this manuscript by Evan DeLucia, Rich Norby, and two anonymous reviewers. Research was supported in part by the US DOE (DE-FG03-00ER63049, DE-FG02-95ER62127) and the US Interagency Terrestrial Ecology (TECO) Initiative, with additional support to RSN from the Nevada Agricultural Experiment Station (Publication no. 52031281) and to SDS from the NSF (DEB-0212812).

References

- Ainsworth EA, Davey PA, Hymus GJ, Osborne CP, Rogers A, Blum H, Nösberger J, Long SP. 2003. Is stimulation of leaf photosynthesis by elevated carbon dioxide concentration maintained in the long term? A test with *Lolium perenne* grown for 10 years at two nitrogen fertilization levels under Free Air CO₂ Enrichment (FACE). *Plant, Cell & Environment* **26**: 705–714.
- Allen AS, Andrews JA, Finzi AC, Matamala R, Richter DD, Schlesinger WH. 2000. Effects of free-air CO₂ enrichment (FACE) on belowground processes in a *Pinus taeda* forest. *Ecological Applications* **10**: 437–448.
- Allen LH Jr. 1992. Free-air CO₂ enrichment experiments: an historical overview. *Critical Reviews in Plant Sciences* **11**: 121–134.
- Amthor JS, Koch GW. 1996. Biotic growth factor β : stimulation of terrestrial ecosystem net primary production by elevated atmospheric CO₂. In: Koch GW, Mooney HA, eds. *Carbon dioxide and terrestrial ecosystems*. San Diego, CA, USA: Academic Press, 399–414.
- Bacastow R, Keeling CD. 1973. Atmospheric carbon dioxide and radiocarbon in the natural carbon cycle. II. Changes from AD 1700–2070 as deduced from a geochemical model. In: Woodwell GM, Pecan EV, eds. *Carbon and the biosphere*. Springfield, VA, USA: US Atomic Energy Commission, National Technical Information Service, 86–135.
- Bazzaz FA. 1990. The response of natural ecosystems to the rising global CO₂ levels. *Annual Review of Ecology and Systematics* **21**: 167–196.
- Belote RT, Weltzin JF, Norby RJ. 2003. Response of an understory plant community to elevated [CO₂] depends on differential responses of dominant invasive species and is mediated by soil water availability. *New Phytologist* **161**: 827–835.
- Bernacchi C, Calfapietra C, Davey PA, Wittig VA, Scarascia-Mugnozza G, Raines CA, Long SP. 2003. Photosynthesis and stomatal conductance responses of poplars to free-air CO₂ enrichment (PopFACE) during the first growth cycle and immediately following coppice. *New Phytologist* **159**: 609–621.
- Billings SA, Schaeffer SM, Zitzer S, Charlet T, Smith SD, Evans RD. 2002. Alterations of nitrogen dynamics under elevated carbon dioxide in an intact Mojave Desert ecosystem: evidence from nitrogen-15 natural abundance. *Oecologia* **131**: 463–467.
- Böhm W. 1979. *Methods of studying root systems*. *Ecological studies* **33**. Berlin Heidelberg, Germany/New York, NY, USA: Springer.
- Bolker BM, Pacala SW, Bazzaz FA, Canham CD, Levin SA. 1995. Species diversity and ecosystem response to carbon dioxide fertilization: conclusions from a temperate forest model. *Global Change Biology* **1**: 373–381.

- Byrne C, Jones MB. 2002. Effects of elevated CO₂ and nitrogen fertiliser on biomass productivity, community structure and species diversity of a semi-natural grassland in Ireland. *Biology and Environment: Proceedings of the Royal Irish Academy* 102B: 141–150.
- von Caemmerer S, Ghannoum O, Conroy JP, Clark H, Newton PCD. 2001. Photosynthetic responses of temperate species to free air CO₂ enrichment (FACE) in a grazed New Zealand pasture. *Australian Journal of Plant Physiology* 28: 439–450.
- Ceulemans R, Mousseau M. 1994. Effects of elevated atmospheric CO₂ on woody plants. *New Phytologist* 127: 425–446.
- Coleman JS, Bazzaz FA. 1992. Effects of CO₂ and temperature on growth and resource use of co-occurring C₃ and C₄ annuals. *Ecology* 73: 1244–1259.
- Craine JM, Reich PB, Tilman D, Ellsworth D, Fargione J, Knops J, Naemm S. 2003. The role of plant species in biomass production and response to elevated CO₂ and N. *Ecology Letters* 6: 623–630.
- Crous KY, Ellsworth DS. 2004. Canopy position affects photosynthetic adjustments to long-term CO₂ enrichment (FACE) in aging needles in a mature *Pinus taeda* L. forest. *Tree Physiology* (In press.)
- Cure JD, Acock B. 1986. Crop responses to carbon dioxide doubling: a literature survey. *Agricultural and Forest Meteorology* 38: 127–145.
- Curtis PS. 1996. A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide. *Plant, Cell & Environment* 19: 127–137.
- Curtis PS, Wang XZ. 1998. A meta-analysis of elevated CO₂ effects on woody plant mass, form, and physiology. *Oecologia* 113: 299–313.
- Daepf M, Suter D, Almeida JPF, Isopp H, Hartwig UA, Frehner M, Blum H, Nösberger J, Lüscher A. 2000. Yield response of *Lolium perenne* swards to free air CO₂ enrichment increased over six years in a high N input system on fertile soil. *Global Change Biology* 6: 805–816.
- DeFalco LA. 2003. Physiological ecology of the invasive annual grass, *Bromus madriensis* subsp. *rubens*, and its interaction with native Mojave Desert species. Dissertation. Reno, NV, USA: University of Nevada-Reno.
- DeLucia EH, Hamilton JG, Naidu SL, Thomas RB, Andrews JA, Finzi A, Lavine M, Matamala R, Mohan JE, Hendrey GR, Schlesinger WH. 1999. Net primary production of a forest ecosystem with experimental CO₂ enrichment. *Science* 284: 1177–1179.
- Díaz S, Grime JP, Harris J, McPherson E. 1993. Evidence of a feedback mechanism limiting plant response to elevated carbon dioxide. *Nature* 364: 616–617.
- Drake BG. 1992. A field study of the effects of elevated CO₂ on ecosystem processes in a Chesapeake Bay wetland. *Australian Journal of Botany* 40: 579–595.
- Drake BG, Leadley PW. 1991. Canopy photosynthesis of crops and native plant communities exposed to long-term elevated CO₂. *Plant, Cell & Environment* 14: 853–860.
- Dukes JS, Mooney HA. 1999. Does global change increase the success of biological invaders? *Trends in Ecology and Evolution* 14: 135–139.
- Ellsworth DS. 2000. Seasonal CO₂ assimilation and stomatal limitations in a *Pinus taeda* canopy. *Tree Physiology* 20: 435–445.
- Ellsworth DS, Oren R, Huang C, Phillips N, Hendrey GR. 1995. Leaf and canopy responses to elevated CO₂ in a pine forest under free-air CO₂ enrichment. *Oecologia* 104: 139–146.
- Evans RD, Rimer R, Sperry L, Belnap J. 2001. Exotic plant invasion alters nitrogen dynamics in an arid grassland. *Ecological Applications* 11: 1301–1310.
- Fajer ED, Bowers MD, Bazzaz FA. 1991. Performance and allocation patterns of the perennial herb, *Plantago lanceolata*, in response to simulated herbivory and elevated CO₂ environments. *Oecologia* 87: 37–42.
- Fajer ED, Bowers MD, Bazzaz FA. 1992. The effect of nutrients and enriched CO₂ environments on production of carbon-based allelochemicals in *Plantago*: a test of the carbon/nutrient balance hypothesis. *American Naturalist* 140: 707–723.
- Farnsworth EJ, Bazzaz FA. 1995. Inter- and intra-generic differences in growth, reproduction, and fitness of nine herbaceous annual species grown in elevated CO₂ environments. *Oecologia* 104: 454–466.
- Farquhar GD, von Caemmerer S, Berry JA. 1980. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* 149: 78–90.
- Fernandez G. 2003a. *Data mining using SAS applications*. Boca Raton, FL, USA: Chapman & Hall/CRC.
- Fernandez G. 2003b. *QLXQLREPEAT. SAS macro for analyzing two qualitative factors with a repeated measures ANOVA*. Reno, NV USA: Department of Applied Economics and Statistics, MS 204. University of Nevada-Reno (<http://www.ag.unr.edu/gf/>).
- Field CB, Chapin FS III, Matson PA, Mooney HA. 1992. Responses of terrestrial ecosystems to the changing atmosphere: a resource-based approach. *Annual Review of Ecology and Systematics* 23: 201–235.
- Filion M, Dutilleul P, Potvin C. 2000. Optimum experimental design for free-air carbon dioxide enrichment (FACE) studies. *Global Change Biology* 6: 843–854.
- Finzi AC, DeLucia EH, Hamilton JG, Richter DD, Schlesinger WH. 2002. The nitrogen budget of a pine forest under free air CO₂ enrichment. *Oecologia* 132: 567–578.
- Garbutt K, Bazzaz FA. 1984. The effects of elevated CO₂ on plants. III. Flower, fruit and seed production and abortion. *New Phytologist* 98: 433–446.
- Gielen B, Calfapietra C, Sabatti M, Ceulemans R. 2001. Leaf area dynamics in a closed poplar plantation under free-air carbon dioxide enrichment. *Tree Physiology* 21: 1245–1255.
- Griffin KL, Tissue DT, Turnbull MH, Whitehead D. 2000. The onset of photosynthetic acclimation to elevated CO₂ partial pressure in field-grown *Pinus radiata* D. Don after 4 years. *Plant, Cell & Environment* 23: 1089–1098.
- Grünzweig JM, Körner C. 2001. Biodiversity effects of elevated CO₂ in species-rich model communities from the semi-arid Negev of Israel. *Oikos* 95: 112–124.
- Gunderson CA, Sholtis JD, Wullschlegel SD, Tissue DT, Hanson PJ, Norby RJ. 2002. Environmental and stomatal control of photosynthetic enhancement in the canopy of a sweetgum (*Liquidambar styraciflua* L.) plantation during 3 years of CO₂ enrichment. *Plant, Cell & Environment* 25: 379–393.
- Hamerlynck EP, Huxman TE, Nowak RS, Redar S, Loik ME, Jordan DN, Zitzer SF, Coleman JS, Seemann JR, Smith SD. 2000. Photosynthetic responses of *Larrea tridentata* to a step-increase in atmospheric CO₂ at the Nevada Desert FACE Facility. *Journal of Arid Environments* 44: 425–436.
- Hanson PJ, McRoberts RE, Isebrands JG, Dixon RK. 1987. An optimal sampling strategy for determining CO₂ exchange-rate as a function of photosynthetic photon flux-density. *Photosynthetica* 21: 98–101.
- Hanson PJ, Todd DE Jr, Amthor JS. 2001. A six-year study of sapling and large-tree growth and mortality responses to natural and induced variability in precipitation and throughfall. *Tree Physiology* 21: 345–358.
- Hättenschwiler S, Handa IT, Egli L, Asshoff R, Ammann W, Körner C. 2002. Atmospheric CO₂ enrichment of alpine treeline conifers. *New Phytologist* 156: 363–375.
- Hebeisen T, Lüscher A, Zanetti S, Fischer BU, Hartwig UA, Frehner M, Hendrey GR, Blum H, Nösberger J. 1997. Growth response of *Trifolium repens* L. and *Lolium perenne* L. as monocultures and bi-species mixture to free air CO₂ enrichment and management. *Global Change Biology* 3: 149–160.
- Heijmans MMPD, Berendse F, Arp WJ, Masselink AK, Klees H, de Visser W, van Breemen N. 2001. Effects of elevated carbon dioxide and increased nitrogen deposition on bog vegetation in the Netherlands. *Journal of Ecology* 89: 268–279.
- Hendrey GR, Kimball BA. 1994. The FACE program. *Agricultural and Forest Meteorology* 70: 3–14.
- Hendrey GR, Ellsworth DS, Lewin KF, Nagy J. 1999. A free air enrichment system for exposing tall forest vegetation to elevated atmospheric CO₂. *Global Change Biology* 5: 293–309.

- Herrick JD, Thomas RB. 2001. No photosynthetic down-regulation in sweetgum trees (*Liquidambar styraciflua* L.) after three years of CO₂ enrichment at the Duke Forest FACE experiment. *Plant, Cell & Environment* 24: 53–64.
- Hooper DU, Vitousek PM. 1997. The effects of plant composition and diversity on ecosystem processes. *Science* 277: 1302–1305.
- Hoosbeek MR, van Breemen N, Berendse F, Grosvernier P, Vasander H, Wallén B. 2001. Limited effect of increased atmospheric CO₂ concentration on ombrotrophic bog vegetation. *New Phytologist* 150: 459–463.
- Houghton JT, Ding Y, Griggs DJ, Noguer M, van der Linden PJ, Xiaosu D, eds. 2001. *Climate Change 2001. The Scientific basis*. Cambridge, UK: Cambridge University Press.
- Housman DC. 2002. Effects of elevated CO₂ on primary productivity in a Mojave Desert ecosystem. Dissertation. Las Vegas, NV, USA: University of Nevada-Las Vegas.
- Huxman TE, Hamerlynck EP, Moore BD, Smith SD, Jordan DN, Zitzer SF, Nowak RS, Coleman JS, Seemann JR. 1998. Photosynthetic down-regulation in *Larrea tridentata* exposed to elevated atmospheric CO₂: interaction with drought under glasshouse and field (FACE) exposure. *Plant, Cell & Environment* 21: 1153–1161.
- Isebrands JG, McDonald EP, Kruger E, Hendrey G, Percy K, Pregitzer K, Sober J, Karnosky DF. 2001. Growth responses of *Populus tremuloides* clones to interacting elevated carbon dioxide and tropospheric ozone. *Environmental Pollution* 115: 359–371.
- Jablonski LM. 1997. Responses of vegetative and reproductive traits to elevated CO₂ and nitrogen in *Raphanus* varieties. *Canadian Journal of Botany* 75: 533–545.
- Jasienski M, Bazzaz FA. 1999. The fallacy of ratios and the testability of models in biology. *Oikos* 84: 321–326.
- Joel G, Chapin FS III, Chiariello NR, Thayer SS, Field CB. 2001. Species-specific responses of plant communities to altered carbon and nutrient availability. *Global Change Biology* 7: 435–450.
- Jones TH, Thompson LJ. 1998. Impacts of rising atmospheric carbon dioxide on model terrestrial ecosystems. *Science* 280: 441–442.
- Keeling CD, Whorf TP. 2002. Atmospheric CO₂ records from sites in the SIO air sampling network. In: *Trends: a compendium of data on global change*. Oak Ridge, TN, USA: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, (<http://cdiac.ornl.gov/trends/co2/sio-mlo.htm>).
- Kimball BA. 1983. Carbon dioxide and agricultural yield: an assemblage and analysis of 430 prior observations. *Agronomy Journal* 75: 779–788.
- Kimball BA, Kobayashi K, Bindi M. 2002. Responses of agricultural crops to free-air CO₂ enrichment. *Advances in Agronomy* 77: 293–368.
- King JS, Pregitzer KS, Zak DR, Sober J, Isebrands JG, Dickson RE, Hendrey GR, Karnosky DF. 2001. Fine-root biomass and fluxes of soil carbon in young stands of paper birch and trembling aspen as affected by elevated atmospheric CO₂ and tropospheric O₃. *Oecologia* 128: 237–250.
- Kopper BJ, Lindroth RL. 2003. Effects of elevated carbon dioxide and ozone on the phytochemistry of aspen and performance of an herbivore. *Oecologia* 134: 95–103.
- Körner C. 1995. Towards a better experimental basis for upscaling plant responses to elevated CO₂ and climate warming. *Plant, Cell & Environment* 18: 1101–1110.
- Körner C. 2000. Biosphere responses to CO₂ enrichment. *Ecological Applications* 10: 1590–1619.
- Kramer PJ. 1981. Carbon dioxide concentration, photosynthesis, and dry matter production. *Bioscience* 31: 29–32.
- Leadley PW, Drake BG. 1993. Open top chambers for exposing plant canopies to elevated CO₂ concentration and for measuring net gas exchange. *Vegetatio* 104/105: 3–15.
- Leadley PW, Niklaus PA, Stocker R, Körner Ch. 1997. Screen-aided CO₂ control (SACC): a middle ground between FACE and open-top chambers. *Acta Oecologia* 18: 225–261.
- Leadley PW, Niklaus PA, Stocker R, Körner C. 1999. A field study of the effects of elevated CO₂ on plant biomass and community structure in a calcareous grassland. *Oecologia* 118: 39–49.
- Lee TD, Tjoelker MG, Ellsworth DS, Reich PB. 2001. Photosynthetic acclimatization of 13 prairie grassland species in the field under elevated carbon dioxide and increased nitrogen. *New Phytologist* 150: 405–418.
- Lee TD, Reich PB, Tjoelker MG. 2003. Legume presence increases photosynthesis and N concentrations of co-occurring non-fixers but does not modulate their responsiveness to carbon dioxide enrichment. *Oecologia* 137: 22–31.
- Leuning R. 1997. Scaling to a common temperature improves the correlation between the photosynthesis parameters J(max) and V-cmax. *Journal of Experimental Botany* 48: 345–347.
- Lewin KF, Hendrey GR, Nagy J, LaMorte RL. 1994. Design and application of a free-air carbon dioxide enrichment facility. *Agricultural and Forest Meteorology* 70: 15–29.
- Loreau M. 2000. Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* 91: 3–17.
- Luo Y, Mooney HA. 1996. Stimulation of global photosynthetic carbon influx by an increase in atmospheric carbon dioxide concentration. In: Koch GW, Mooney HA, eds. *Carbon dioxide and terrestrial ecosystems*. San Diego, CA, USA: Academic Press, 381–397.
- Luo Y, Reynolds JF. 1999. Validity of extrapolating field CO₂ experiments to predict carbon sequestration in natural ecosystems. *Ecology* 80: 1568–1583.
- Luo Y, Reynolds JF, Wang Y, Wolfe D. 1999. A search for predictive understanding of plant responses to elevated [CO₂]. *Global Change Biology* 5: 143–156.
- Luumala EM, Laitinen K, Kellomäki S, Vapaavuori E. 2003. Variable photosynthetic acclimation in consecutive cohorts of Scots pine needles during 3 years of growth at elevated CO₂ and elevated temperature. *Plant, Cell & Environment* 26: 645–660.
- Lüscher A, Hendrey GR, Nösberger J. 1998. Long-term responsiveness to free air CO₂ enrichment of functional types, species and genotypes of plants from fertile permanent grassland. *Oecologia* 113: 37–45.
- Matamala R, Schlesinger WH. 2001. Effects of elevated atmospheric CO₂ on fine root production and activity in an intact temperate forest ecosystem. *Global Change Biology* 6: 967–979.
- McConaughay KDM, Berntson GM, Bazzaz FA. 1993. Limitations to CO₂-induced growth enhancement in pot studies. *Oecologia* 94: 550–557.
- McLeod AR, Long SP. 1999. Free-air carbon dioxide enrichment (FACE) in global change research: a review. *Advances in Ecological Research* 28: 1–56.
- Medlyn BE, Badeck F-W, De Pury DGG, Barton CVM, Broadmeadow M, Ceulemans R, De Angelis P, Forstreuter M, Jach ME, Kellomäki S, Laitat E, Marek M, Philippot S, Rey A, Strassmeyer J, Laitinen K, Liozon R, Portier B, Roberntz P, Wang K, Jarvis PG. 1999. Effects of elevated [CO₂] on photosynthesis in European forest species: a meta-analysis of model parameters. *Plant, Cell & Environment* 22: 1475–1495.
- Medlyn BE, Barton CVM, Broadmeadow MSJ, Ceulemans R, De Angelis P, Forstreuter M, Freeman M, Jackson SB, Kellomäki S, Laitat E, Rey A, Roberntz P, Sigurdsson BD, Strassmeyer J, Wang K, Curtis PS, Jarvis PG. 2001. Stomatal conductance of forest species after long-term exposure to elevated CO₂ concentration: a synthesis. *New Phytologist* 149: 247–264.
- Melillo JM, Janetos AC, Karl TR, Barron EJ, Burkett V, Cecich TF, Corell R, Jacobs K, Joyce L, Miller B, Morgan MG, Parson EA, Richels RG, Schimel DS. 2001. *Climate change impacts on the United States: the potential consequences of climate variability and change*. Cambridge, UK: Cambridge University Press.
- Melillo JM, McGuire AD, Kicklighter DW, Moore B III, Vorosmarty CJ, Schloss AL. 1993. Global climate change and terrestrial net primary production. *Nature* 363: 234–240.

- Miglietta F, Hoosbeek MR, Foot J, Gigon F, Hassinen A, Heijmans M, Peressotti A, Saarinen T, Van Breemen N, Wallén B. 2001a. Spatial and temporal performance of a miniFACE (Free Air CO₂ Enrichment) system on bog ecosystems in northern and central Europe. *Environmental Monitoring and Assessment* 66: 107–127.
- Miglietta F, Peressotti A, Vaccari FP, Zaldei A, De Angelis O, Scarascia-Mugnozza G. 2001b. Free-air CO₂ enrichment (FACE) of a poplar plantation: the POPFACE fumigation system. *New Phytologist* 150: 465–476.
- Mitchell CE, Reich PB, Tilman D, Groth JV. 2003. Effects of elevated CO₂, nitrogen deposition, and decreased species diversity on foliar fungal plant disease. *Global Change Biology* 9: 438–451.
- Mooney HA, Koch GW. 1994. The impact of rising CO₂ concentrations on the terrestrial biosphere. *Ambio* 23: 74–76.
- Morgan JA, Mosier AR, Milchunas DG, LeCain DR, Nelson JA, Parton WJ. 2004a. CO₂ enhances productivity, alters species composition, and reduces forage digestibility of shortgrass steppe vegetation. *Ecological Applications* 14: 208–219.
- Morgan JA, Pataki DE, Körner C, Clark H, Del Grosso SJ, Grünzweig JM, Knapp AK, Mosier AR, Newton PCD, Niklaus PA, Nippert J, Nowak RS, Parton WJ, Polley HW, Shaw MR. 2004b. Water relations in grassland and desert ecosystems exposed to elevated atmospheric CO₂. *Oecologia* (In press.)
- Morison JIL. 1985. Sensitivity of stomata and water use efficiency to high CO₂. *Plant, Cell & Environment* 8: 467–474.
- Myers DA, Thomas RB, DeLucia EH. 1999. Photosynthetic capacity of loblolly pine (*Pinus taeda* L.) trees during the first year of carbon dioxide enrichment in a forest ecosystem. *Plant, Cell & Environment* 22: 473–481.
- Naumburg E, Housman DC, Huxman TE, Charlet TN, Loik MI, Smith SD. 2003. Photosynthetic responses of Mojave Desert shrubs to free air CO₂ enrichment are greatest during wet years. *Global Change Biology* 9: 276–285.
- Neilson RP. 1995. A model for predicting continental-scale vegetation distribution and water balance. *Ecological Applications* 5: 362–385.
- Niklaus PA, Leadley PW, Schmid B, Körner Ch. 2001. A long-term field study on biodiversity–elevated CO₂ interactions in grassland. *Ecological Monographs* 71: 341–356.
- Niklaus PA, Alpehi J, Ebersberger D, Kampichler C, Kandeler E, Tschirko D. 2003. Six years of *in situ* CO₂ enrichment evoke changes in soil structure and soil biota of nutrient-poor grassland. *Global Change Biology* 9: 585–600.
- Noormets A, Sober A, Karnosky DF. 2001. Stomatal and non-stomatal limitation to photosynthesis in two trembling aspen (*Populus tremuloides* Michx.) clones exposed to elevated CO₂ and/or O₃. *Plant, Cell & Environment* 24: 327–336.
- Norby RJ, Wullschlegel SD, Gunderson CA. 1996. Tree responses to elevated CO₂ and implications for forests. In: Koch GW, Mooney HA, eds. *Carbon dioxide and terrestrial ecosystems*. San Diego, CA, USA: Academic Press, 1–21.
- Norby RJ, Wullschlegel SD, Gunderson CA, Johnson DW, Ceulemans R. 1999. Tree responses to rising CO₂ in field experiments: implications for the future forest. *Plant, Cell & Environment* 22: 683–714.
- Norby RJ, Kobayashi K, Kimball BA. 2001. Rising CO₂ – future ecosystems. *New Phytologist* 150: 215–221.
- Norby RJ, Hanson PJ, O'Neill EG, Tschaplinski TJ, Weltzin JF, Hansen RA, Cheng W, Wullschlegel SD, Gunderson CA, Edwards NT, Johnson DW. 2002. Net primary productivity of a CO₂-enriched deciduous forest and the implications for carbon storage. *Ecological Applications* 12: 1261–1266.
- Nowak RS, Jordan DN, DeFalco LA, Wilcox CS, Coleman JS, Seemann JR, Smith SD. 2001. Leaf conductance decreased under free-air CO₂ enrichment (FACE) for three perennials in the Nevada desert. *New Phytologist* 150: 449–458.
- Ollinger SV, Aber JD, Reich PB, Freuder RJ. 2002. Interactive effects of nitrogen deposition, tropospheric ozone, elevated CO₂ and land use history on the carbon dynamics of northern hardwood forests. *Global Change Biology* 8: 545–562.
- Oren R, Ellsworth DS, Johnsen KH, Phillips N, Ewers BE, Maler C, Schäfer KVR, McCarthy H, Hendrey G, McNulty SG, Katul GG. 2001. Soil fertility limits carbon sequestration by forest ecosystems in a CO₂-enriched atmosphere. *Nature* 411: 469–472.
- Owensby CE, Ham JM, Knapp AK, Auen LA. 1999. Biomass production and species composition change in a tallgrass prairie ecosystem after long-term exposure to elevated atmospheric CO₂. *Global Change Biology* 5: 497–506.
- Pearcy RW, Björkman O. 1983. Physiological effects. In: Lemon E, ed. *CO₂ and plants: the response of plants to rising levels of atmospheric carbon dioxide*. AAAS selected symposium 84. Washington, DC, USA: AAAS, 65–105.
- Pepin S, Körner C. 2002. Web-FACE: a new canopy free-air CO₂ enrichment system for tall trees in mature forests. *Oecologia* 133: 1–9.
- Percy KE, Awmack CS, Lindroth RL, Kubiske ME, Kopper BJ, Isebrands JG, Pregitzer KS, Hendrey GR, Dickson RE, Zak DR, Okanen E, Sober J, Harrington R, Karnosky DF. 2002. Altered performance of forest pests under atmospheres enriched by CO₂ and O₃. *Nature* 420: 403–407.
- Peterson AG, Ball JT, Luo YQ, Field CB, Reich PB, Curtis PS, Griffin KL, Gunderson CA, Norby RJ, Tissue DT, Forstreuter M, Rey A, Vogel CS. 1999. The photosynthesis leaf nitrogen relationship at ambient and elevated atmospheric carbon dioxide: a meta-analysis. *Global Change Biology* 5: 331–346.
- Phillips DL, Tingey DT, Johnson MG, Catricala CE, Hoyman TL. 2002. *Effects of elevated CO₂ on fine root dynamics in a Mojave Desert ecosystem: a FACE study*. Abstracts, Ecological Society of America 87th annual meeting. Washington, DC, USA: Ecological Society of America, 409.
- Pinter PJ Jr, Kimball BA, Wall GW, LaMorte RL, Hunsaker DJ, Adamsen FJ, Frumau KFA, Vufts HF, Hendrey GR, Lewin KF, Nagy J, Johnson HB, Wechsung F, Leavitt SW, Thompson TL, Matthias AD, Brooks TJ. 2000. Free-air CO₂ enrichment (FACE): blower effects on wheat canopy microclimate and plant development. *Agricultural and Forest Meteorology* 103: 319–333.
- Poorter H, Navas M-L. 2003. *Tansley review: plant growth and competition at elevated CO₂: on winners, losers and functional groups*. *New Phytologist* 157: 175–198.
- Pritchard SG, Roger HH, Davis MA, van Santen E, Prior SA, Schlesinger WH. 2001. The influence of elevated atmospheric CO₂ on fine root dynamics in an intact temperate forest. *Global Change Biology* 7: 829–837.
- Reich PB, Ellsworth DS, Walters MB, Vose JM, Gresham C, Volin JC, Bowman WD. 1998. Generality of leaf trait relationships: a test across six biomes. *Ecology* 80: 1955–1969.
- Reich PB, Knops J, Tilman D, Craine J, Ellsworth D, Tjoelker M, Lee T, Wedin D, Naeem S, Bahauddin D, Hendrey G, Jose S, Wrage K, Goth J, Bengston W. 2001a. Plant diversity enhances ecosystem responses to elevated CO₂ and nitrogen deposition. *Nature* 410: 809–812.
- Reich PB, Tilman D, Craine J, Tjoelker MG, Knops J, Wedin D, Naeem S, Bahauddin D, Goth J, Bengston W, Lee TD. 2001b. Do species and functional groups differ in acquisition and use of C, N and water under varying atmospheric CO₂ and N availability regimes? A field test with 16 grassland species. *New Phytologist* 150: 435–448.
- Rogers A, Ellsworth DS. 2002. Photosynthetic acclimation of *Pinus taeda* (loblolly pine) to long-term growth in elevated pCO₂ (FACE). *Plant, Cell & Environment* 25: 851–858.
- Sage RF. 1994. Acclimation of photosynthesis to increasing atmospheric CO₂: the gas exchange perspective. *Photosynthesis Research* 39: 351–368.
- Sage RF. 1996. Modification of fire disturbance by elevated CO₂. In: Körner C, Bazzaz FA, eds. *Carbon dioxide, populations, and communities*. San Diego, CA, USA: Academic Press, 231–249.

- SAS. 2001. *SAS system for Windows, v8.02*. Cary, NC, USA: SAS Institute.
- Saxe H, Ellsworth DS, Heath J. 1998. Tree and forest functioning in an enriched CO₂ atmosphere. *New Phytologist* 139: 395–436.
- Schäfer KVR, Oren R, Lai C-T, Katul GG. 2002. Hydrologic balance in an intact temperate forest ecosystem under ambient and elevated atmospheric CO₂ concentration. *Global Change Biology* 8: 895–911.
- Schortemeyer M, Hartwig U, Hendrey G, Sadowsky MJ. 1996. Microbial community changes in the rhizospheres of white clover and perennial ryegrass exposed to free air carbon dioxide enrichment (FACE). *Soil Biology and Biochemistry* 28: 1717–1724.
- Shaw MR, Zavaleta ES, Chiariello NR, Cleland EE, Mooney HA, Field CB. 2002. Grassland responses to global environmental changes suppressed by elevated CO₂. *Science* 298: 1987–1990.
- Smith SD, Huxman TE, Zitzer SF, Charlet TN, Housman DC, Coleman JS, Fenstermaker LK, Seemann JR, Nowak RS. 2000. Elevated CO₂ increases productivity and invasive species success in an arid ecosystem. *Nature* 408: 79–82.
- Stitt M, Krapp A. 1999. The interaction between elevated carbon dioxide and nitrogen nutrition: the physiological and molecular background. *Plant, Cell & Environment* 22: 583–621.
- Strain BR. 1987. Direct effects of increasing atmospheric CO₂ on plants and ecosystems. *Trends in Ecology and Evolution* 2: 18–21.
- Strain BR, Bazzaz FA. 1983. Terrestrial plant communities. In: Lemon E, ed. *CO₂ and plants: the response of plants to rising levels of atmospheric carbon dioxide*. AAAS Selected Symposium 84. Washington, DC, USA: AAAS, 177–222.
- Takeuchi Y, Kubiske ME, Isebrands JG, Pregitzer KS, Hendrey G, Karnosky DF. 2001. Photosynthesis, light and nitrogen relationships in a young deciduous forest canopy under open-air CO₂ enrichment. *Plant, Cell & Environment* 24: 1257–1268.
- Tilman D, Knops J, Wedin D, Reich P, Ritchie M, Siemann E. 1997. The influence of functional diversity and composition on ecosystem processes. *Science* 277: 1300–1302.
- Tissue DT, Megonigal JP, Thomas RB. 1997. Nitrogenase activity and N₂ fixation are stimulated by elevated CO₂ in a tropical N₂-fixing tree. *Oecologia* 109: 28–33.
- Wand SJE, Midgley GF, Jones MH, Curtis PS. 1999. Responses of wild C₄ and C₃ grass (Poaceae) species to elevated atmospheric CO₂ concentration: a test of current theories and perceptions. *Global Change Biology* 5: 723–740.
- Watson RT, Zinyowera MC, Moss RH. 1998. *The regional impacts of climate change: an assessment of vulnerability*. Cambridge, UK: Cambridge University Press.
- Weltzin JF, Belote RT, Sanders NJ. 2003. Biological invaders in a greenhouse world: will elevated CO₂ fuel plant invasions? *Frontiers in Ecology and the Environment* 1: 146–153.
- Whittaker RH. 1970. *Communities and ecosystems*. New York, NY, USA: MacMillan.
- Wullschleger SD, Norby RJ, Gunderson CA. 1997. Forest trees and their response to atmospheric carbon dioxide enrichment: a compilation of results. In: *Advances in carbon dioxide effects research. ASA special publication no. 61*. Madison, WI, USA: ASA, 79–100.
- Yin X. 2002. Responses of leaf nitrogen concentration and specific leaf area to atmospheric CO₂ enrichment: a retrospective analysis. *Global Change Biology* 8: 631–642.
- Zak DR, Pregitzer KS, Curtis PS, Teeri JA, Fogel R, Randlett DL. 1993. Elevated atmospheric CO₂ and feedback between carbon and nitrogen cycles. *Plant and Soil* 151: 105–117.
- Zak DR, Pregitzer KS, King JS, Holmes WE. 2000. Elevated atmospheric CO₂, fine roots and the response of soil microorganisms: a review and hypothesis. *New Phytologist* 147: 201–222.
- Zak DR, Holmes WE, Finzi AC, Norby RJ, Schlesinger WH. 2003. Soil nitrogen cycling under elevated CO₂: a synthesis of forest FACE experiments. *Ecological Applications* 13: 1508–1514.
- Zavaleta ES. 2001. Influences of climate and atmospheric changes on plant diversity and ecosystem function in a California grassland. PhD Dissertation. Stanford, CA, USA: Stanford University.
- Zavaleta ES, Shaw MR, Chiariello NR, Mooney HA, Field CB. 2003. Additive effects of simulated climate changes, elevated CO₂, and nitrogen deposition on grassland diversity. *Proceedings of the National Academy of the USA* 100: 7650–7654.
- Ziska LH. 2003. Evaluation of the growth response of six invasive species to past, present and future atmospheric carbon dioxide. *Journal of Experimental Botany* 54: 395–404.



About New Phytologist

- *New Phytologist* is owned by a non-profit-making **charitable trust** dedicated to the promotion of plant science, facilitating projects from symposia to open access for our Tansley reviews. Complete information is available at www.newphytologist.org
- Regular papers, Letters, Research reviews, Rapid reports and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as-ready' via *OnlineEarly* – average first decisions are just 5–6 weeks. Essential colour costs are **free**, and we provide 25 offprints as well as a PDF (i.e. an electronic version) for each article.
- For online summaries and ToC alerts, go to the website and click on 'Journal online'. You can take out a **personal subscription** to the journal for a fraction of the institutional price. Rates start at £108 in Europe/\$193 in the USA & Canada for the online edition (click on 'Subscribe' at the website)
- If you have any questions, do get in touch with Central Office (newphytol@lancaster.ac.uk; tel +44 1524 592918) or, for a local contact in North America, the USA Office (newphytol@ornl.gov; tel 865 576 5261)